

[DH and HPA logos]

**Health Effects of Climate Change in the
UK**

**An update of the Department of Health
report
2001/2002**

DRAFT FOR COMMENT

Department of Health

and

Health Protection Agency

2007

DH INFORMATION READER BOX

Policy	Estates
HR / Workforce	Performance
Management	IM & T
Planning	Finance
Clinical	Partnership Working

Document Purpose	Consultation/Discussion
ROCR Ref:	Gateway Ref: 8014
Title	Health Effects of Climate Change in the UK
Author	DH / Health Protection Agency
Publication Date	04 May 2007
Target Audience	Climate change scientists
Circulation List	Voluntary Organisations/NDPBs
Description	A DH commissioned independent scientific report updating earlier DH report on health effects of climate change. Report to be placed on web for comment. Comments to be submitted to I. Myers in writing, either via e-mail or hard copy within a 6 week period.
Cross Ref	Health Effects of Climate Change in the UK 2001
Superseded Docs	N/A
Action Required	N/A
Timing	By 08 Jun 2007
Contact Details	Isabella Myers Air Pollution Unit - CHaPD Health Protection Agency Chilton, Didcot OX11 0RQ 07881 512077 www.dh.gov.uk/airpollution www.hpa.org.uk
For Recipient's Use	

Preface I

Climate change will have consequences for the health of UK citizens. In 2002 the Department of Health published a report on the '*Health Effects of Climate Change in the UK*'. This report was amongst the first of its kind in that it sought to provide quantitative estimates of the possible impacts of climate change on health. The report was well received and widely quoted.

Now, five years later, I am pleased to welcome an update to the original report, published jointly by the DH and the Health Protection Agency. Independent scientific experts were commissioned by DH to focus on areas that had changed since the original report was written. The briefer new report is therefore to be read in conjunction with the previous report. It is being published as a draft for wider comment.

Since the work of this expert panel, other key reports on climate change have been published, most recently that of the Intergovernmental Panel on Climate Change (IPCC). Our report supports the IPCC findings but concentrates on the impacts of climate change on human health rather than the factors which affect climate change itself.

Actions are being taken across the UK Government and the NHS to reduce emissions contributing to climate change. The UK's Climate Change Bill is the first legislation of its kind in the world, establishing a long-term legal framework to underpin the UK's contribution to tackling climate change, by putting in place a clear and credible emissions reduction pathway to a statutory goal of a 60% reduction in carbon dioxide emissions by 2050.

From the Government's Climate Change programme DH is leading in setting mandatory targets for energy/carbon efficiency for the public sector and is working with the Carbon Trust on the NHS Carbon Management Programme. Work is continuing across the NHS at a regional and local level to mitigate to the projected impacts of climate change. The NHS is working towards mandatory energy and carbon efficiency targets (from 2000 to 2010) with advice and guidance from DH on energy/ carbon management for existing operational estate and for new capital build developments. The energy performance of the NHS is improving, with over 70% meeting the mandatory targets of 55-65 gigajoules/100m³.

In January 2007 Andy Burnham announced a £100M Energy and Sustainability Capital Fund for 07-March 09 to assist the NHS to meet the overarching target of 15% energy or 0.15 million tonnes carbon efficiency saving between 2000 to 2010, particularly by spend to save initiatives.

DH is continuing to work with the Sustainable Development Commission on the NHS as a Corporate Citizen self assessment model already taken up by 151 NHS bodies which includes action to make operations more sustainable. The Government SDIG programme requires reporting on sustainable developments in government and addressing operational targets on energy use, water and waste.

One of the effects of climate change already encountered in this country is the increased frequency of heatwaves. The devastating heatwave across Europe in 2003 led to DH first launching its *National Heatwave Plan* in 2004, in which a 'Heat-Health Watch' system operates in England during the summer months, with advice from the Met Office, with four levels of response and appropriate advice.

Key areas for the NHS in adapting to climate change include: adapting the health and social care infrastructure (hospitals, nursing homes) to be more resilient to the effects of heat, gales and floods; development of local 'Heatwave', 'Gale' and 'Flood' plans for coping with disasters; and increasing awareness of how people can adapt to changes in climate.

Climate change poses great challenges and it is important to plan ahead for the health consequences. Let's not forget we also have a societal role to play in the mitigation of climate change by supporting sustainable development programmes – through consumer choice, reducing our carbon footprints and recycling waste. I hope you will take time to consider this report and let us have your comments.

CAROLINE FLINT

Preface II

Climate change is perhaps the most significant environmental problem which mankind will face in the coming century. Efforts to reduce the extent of climate change are of course important, but it is likely that we will have to deal with at least some impacts on health. Preparing for climate change is now one of the top 4 shared priorities for UK Action set out in 2005 in *Securing the Future, delivering the UK sustainable development strategy*.

As a response to the need to estimate possible impacts of climate change on health, this new report, jointly published by the Department of Health and the Health Protection Agency, is a timely update of earlier work published by the Department of Health in 2002 (Health Effects of Climate Change in the UK).

The new report should be read in conjunction with the original report as it focuses on areas where things have changed rather than covering, again, the whole field. A new approach has been taken in some sections and recent research findings have been included in many chapters.

The report was written by Dr. R.L. Maynard and a group of UK experts and we are grateful to them for their input to this incisive report and for the time they devoted to it.

The report was edited by Sari Kovats. Her contribution has been most important, indeed the report would not have been completed without her input.

Sir William Stewart

Chairman

Health Protection Agency

Reference

HM Government (2005) *Securing the Future, delivering the UK sustainable development strategy*. TSO – The Stationery Office.

http://www.sustainable-development.gov.uk/publications/pdf/strategy/SecFut_complete.pdf

List of Contributors

M. Allen (University of Oxford)

H.R. Anderson (St. George's, University of London)

G. Bentham (School of Environmental Sciences, University of East Anglia)

R.G. Derwent (Rd Scientific)

G.D. Donaldson (Queen Mary, University of London)

C. Goodess (Climatic Research Unit, University of East Anglia)

G.D. Hayman (AEA Technology)

T. Holt (Climatic Research Unit, University of East Anglia)

W.R. Keatinge (Queen Mary, University of London)

S. Kovats (London School of Hygiene and Tropical Medicine)

S. Lindsay (Durham University)

R.L. Maynard (Health Protection Agency)

I. Myers (Department of Health Toxicology Unit, Imperial College)

G. Nichols (Centre for Infections, Health Protection Agency)

S. Randolph (University of Oxford)

D. Rogers (University of Oxford)

J.R. Stedman (AEA Technology)

D. Viner (Climatic Research Unit, University of East Anglia)

S.G. Willis (Durham University)

Executive Summary

The Department of Health report, “Health Effects of Climate Change in the UK”, published in 2002, was amongst the first of its kind in that it focused on the quantitative aspects of possible impacts of climate change on health. The report was well received both in the UK and abroad. The present report provides an update on the findings of the first report but is not a second edition. Much that was included in the original report remains of value and detailed accounts of some approaches to quantification of effects have not been repeated here. In this second report a different approach has been taken in some of the major areas of concern. Preparation for climate change is now one of the top 4 shared priorities for UK Action set out in 2005 in *Securing the Future, delivering the UK sustainable development strategy*.

A small group of experts were asked to consider the original report and to advise on whether updating was needed. The group included chapter authors from the group that wrote the original report and experts who had not been involved before. For some chapters revision was recommended; for others, substantial rewriting was undertaken.

An important feature of the report is the updating of the climate change scenarios provide in Chapter 2. These reflect state of the art work: the UK is amongst world leaders in this area of prediction of climate change. It is now perhaps even more clear than it was in 2002 that the climate of the UK is changing. The latest models used in this report, predict an increase of mean annual temperature in the UK of between 2.5 and 3 degrees Centigrade by the end of the century. Periods of very cold weather will become less common, but periods of very hot weather (heatwaves) will become more common. Application of the epidemiological concept of ‘attributable risk’ to meteorological data allows us to assess the extent to which human influence on climate has contributed to the risk of a specific weather event. Using this approach, a significant role for human influence can already be identified in, for example, the European summer heatwave of 2003 which contributed to over 14,000 premature deaths in France. Changes in wind and rainfall are less certain but periods of sudden heavy rain seem likely to become more common despite a possible reduction in annual rainfall in some areas. Flooding is an increasing risk.

We know rather little of the total effects on health of flooding. These were discussed in some detail in the original report and that discussion has not been repeated here. The predicted risk of severe coastal flooding remains low, but this will increase as sea levels rise. The need for upgraded coastal defences in East Anglia is stressed.

As regards vector-borne diseases the picture is perhaps more encouraging. Reappraisal of the evidence suggests that outbreaks of malaria in the UK are likely to remain rare, though Health Authorities need to remain alert to the possibility of outbreaks of malaria in other European countries and to the possibility that more effective vectors (different species on mosquito) may arrive in the UK. Rapid response to outbreaks of malaria will reduce the chances of the disease becoming endemic in the UK. Tick-borne disease are likely to become more common in the UK, but this will be more likely to be due to changes in land use and leisure activities than to climate change. The likelihood that Tick-borne encephalitis will become established in the UK is very low.

Warmer summers are likely to be associated with an increase in foodborne diseases but the picture has not changed significantly since the original report was published.

A fresh look at the effects of climate change on supplies of drinking water has been taken. Three problems have been identified: increased rainfall (over short periods) leading to increased numbers of bacteria in surface water, increased water temperature leading in an increase in algal blooms in reservoirs and a decrease in the efficiency of chemical coagulation: a major method of removal of microbes from drinking water.

A new approach to the direct effects of high temperatures on health has been taken in this report. This chapter should be read in conjunction with that in the original report: both contain important and complimentary material. Summers in the UK have become warmer, but no change in heat related deaths occurred during the period 1971-2003. This suggests that the UK population is capable of adapting to warmer conditions. But heatwaves still present a serious risk. Predicting severe heatwaves and their effects is difficult, but there is a 1 in 40 chance that by 2012 South-Eastern England will have experienced a severe heatwave that will cause perhaps 3000 immediate heat-related deaths and 6350 heat-related deaths. In terms of conventional thinking about risks to health a risk of 1 in 40 is high. Winter deaths will continue to decline as the climate warms.

The air pollution climate of the UK will continue to change. Though concentrations of a number of important pollutants are likely to decline over the next half century, the concentration of ozone is likely to increase. This will increase attributable deaths and hospital admissions. The increases are likely to be significant: with the least constraining assumptions (no threshold of effect assumed) up to about 1500 extra deaths and hospital admissions per annum might be expected.

Further work has confirmed the effects on increased exposure to ultra-violet light. Skin cancers are expected to increase.

In each chapter of the report summaries of the main points are made and a number of recommendations for mitigating the effects of climate change on health are put forward. These can be summarised as:

- The need for greater emphasis to be placed on climate change and its impacts and the need for governments to focus on this problem.
- Measures individuals can take to mitigate the effects of climate change on their health. Keeping cool in hot weather is important. The easy to remember advice: “keep cool, keep clean, keep covered” remains sensible.
- The need for further research in many of the areas touched on in this report.

Contents

Preface I	i
Preface II	iii
Contributors	iv
Executive summary	v
1 Climate change in UK: the evidence	1
1.1 Introduction.....	2
1.2 Observed trends in UK Climate	2
1.3 Future Climate Change in the UK: the UKCIP02 scenarios.....	5
1.4 Future Climate Change in the UK: extreme events	8
1.5 Changes in the frequency of heatwaves of the order of the 2003 event	18
1.6 Low probability, high impact events	20
2 Flooding and windstorms and climate change	24
2.1 Introduction.....	25
2.2 The health effects of flooding and windstorms.....	25
2.3 Climate change, riverine flooding and extreme precipitation	27
2.4 Climate change and windstorms	30
2.5 Sea level rise, coastal flooding and storm surges.....	31
2.6 Populations at risk of flooding	32
3 Vector-borne diseases and climate change.....	35
3.1 Introduction.....	36
3.2 Methodology	36
3.3 Malaria	39
3.4 Tick-borne infections.....	50
3.5 Possible sources of other vector-borne diseases.....	57
3.5 Other insect-related problems	66
3.6 Priorities for future action	66
4 Foodborne disease and climate change	72
4.1 Introduction.....	73
4.2 Current impacts of climate and weather	73
4.3 Future impacts of climate change	75
5 Water and disease and climate change	77
5.1 Introduction.....	78
5.2 The effects of heavy rainfall on outbreaks of waterborne disease	78
5.3 The effect of drought on water quality and water availability	79
5.4 Recreational waters and algal blooms.....	80
6 Direct effects of rising temperatures on mortality in the UK.....	82
6.1 Introduction.....	83
6.2 Observed changes in heat and cold-related mortality (1971-2003)	83
6.3 Observed and future changes in heatwaves and in heatwave mortality	87
6.4 Discussion.....	89
6.5 Actions to reduce heatwave mortality as summers become hotter.	90
7 The health impact of climate change due to changes in air pollution	92
7.1 Introduction.....	93
7.2 Meteorological factors, emissions, and future urban air pollution episodes.....	93
7.3 Health impacts: Episodes	99
7.4 Quantitative health impact assessment	99
8 Climate change, ground level UV and health.....	107
8.1 Introduction.....	108
8.2 Health impacts of UV	108
8.3 Changing exposures due to changes in behaviour, climate and ozone layer	109
8.4 Future impacts on health.....	109

1 Climate change in UK: the evidence

(See also Chapter 3 in 2001 report)

T Holt, Climatic Research Unit, University of East Anglia

C Goodess, Climatic Research Unit, University of East Anglia

D Viner, Climatic Research Unit, University of East Anglia

Summary

- Mean annual Central England temperatures have continued to rise and are now over 2°C higher than in the coldest period of the 'Little Ice Age' in about 1690. Half of this increase occurred in the last 40 years, which has also seen a rapid rise in the number of hot days. These changes are consistent with anthropogenic warming.
- Total summer rainfall in England and Wales has decreased by about 50 mm over the last 250 years, while autumn and winter rainfall have both increased by about 50 mm.
- Assuming a medium-high emissions scenario, the increase in mean temperature over the UK by the end of the century is projected to be about 2.0° C ($\pm 0.2^{\circ}$ C) in winter and about 4.0° C ($\pm 0.3^{\circ}$ C) in summer.
- The duration of the annual longest cold spell (a period of days with the lowest daily temperature below 0° C) is projected to decrease all over the UK, with the greatest reduction of about 7 days (± 1 day) over Scotland.
- Hot spells (defined as periods with daily maximum temperature higher than 25° C) are projected to be prolonged on average by up to 10 days (+1 to 2 days) over Central and Southern England, and by less than 5 days over the rest of the UK.
- Changes in average annual total rainfall have too much uncertainty to say more than that Scotland is likely to become about 10% drier by the end of the century and most of the rest of the UK will not change very much. The average annual maximum length of drought is projected to extend by about 4 to 6 days (± 1 day) over much of England and Wales.
- Episodes of very high wind speeds are expected to decline over much of the UK under global warming. However, increases are noted over Wales, Northeast England, and Northwest Scotland. Changes in the average annual highest maximum daily wind speed are trivial and uncertain.
- Heatwaves of the order of the 2003 event for England will be encountered only occasionally until about 2030. After this, however, they can be expected much more frequently, becoming much more severe in intensity and duration after 2060.
- A shutdown of the North Atlantic thermohaline circulation system leading to colder winters is considered highly unlikely on the centennial timescale considered here.

Recommendations

- o Construction of refined climate scenarios focused on health impacts. This will require work on the following issues:
 - o Further downscaling to finer spatial scales.
 - o Improved representation of urban areas and their heat sources in climate models, in order to model changes in the urban heat island effect.
 - o Focus on indices of extremes of greatest concern for health impacts, e.g., threshold temperatures used in the Heatwave Plan for England.
 - o Consistency with the UKCIP next scenarios, which will be probabilistic.

1.1 Introduction

Since the first report on the “Health Effects of Climate Change in the UK”, the assessment of future climate change has been updated in the revised national scenarios (UKCIP02). The observed warming trends have been confirmed with more recent years. Globally, 2005 was the second hottest year after 1998.

A new generation of high resolution climate models (Regional Climate Models or RCMs) allow for improved estimates of future changes in the frequency, intensity and duration of extreme events in the UK.

This chapter briefly describes the current scientific consensus on the current and future effects of climate change on the UK climate, as well as introducing the scenarios that are used in the later chapters to estimate future health effects of climate change.

1.2 Observed trends in UK Climate


1.2.1 Observed warming in England

The Central England Temperature series (CET) provides time-series of air temperature at the surface for an area covered by the triangle with coordinates Preston, Bristol, and London. The mean monthly series is available from 1659 to present and is the longest instrumental temperature record in the world (Manley, 1953 and 1974). Daily time-series of the CET are available from 1772, and daily maxima and minima from 1878 (Parker *et al*, 1992).

Figure 1.1 shows the mean annual CET series based on monthly data from 1659 to 2004. The trend is shown using a numerical filter. This decomposes the time-series into its non-stationary time trend and stationary residual components. Here, the term “stationary” is used in its statistical sense to mean a time-series with a constant mean. Clearly, therefore, a time series with trend induced by global warming, for example, must be a non-stationary series. The same filter is applied to all time-series in this chapter.

Following the coldest period of the “Little Ice Age” in about 1690, temperatures in the Midlands climbed until about the year 1730, then fell by 0.5° C until 1750, and remained essentially stable until about 1900. Temperatures then peaked once more around the year 1950, fell slightly until 1960, and then climbed steeply (by about 1° C) to the present day. The detailed inset in Figure 1.1 focuses on the period 1960 to 2004, when temperatures started and continued to rise. Mean annual CET temperatures are now 2° C higher than in the coldest period of the “Little Ice Age” (Figure 1.1). Half of this increase occurred in the last 40 years, and is consistent with theories of anthropogenic warming of the atmosphere. In the case of global temperature, the IPCC has stated that ‘there is new and stronger evidence that most of the warming observed over the last 50 years is attributable to human activities’ (Houghton *et al*, 2001).

Figure 1.1 Mean annual Central England Temperature (CET): full record and the last 44 years



1.2.2 Hot days and days below freezing

The number of hot days in the UK has been increasing along with mean temperature. Both increases are consistent with theories of anthropogenic climate change. The number of days per year when mean daily CET temperatures exceeded 20° C (Figure 1.2a) was quite stable until about 1960, after which temperatures began a persistent and rapid climb to present levels.

The behaviour of the number of days per year with temperatures below 0° C is, however, quite different, with a continuous decrease since the 1770s (Figure 1.2b). The onset of anthropogenic warming coincides with the end of the “Little Ice Age”, so the ambient climate, with respect to very cold days, was considered as warming. It is difficult to disentangle the influence of the two causes. However, it is clear that daily variability has reduced considerably since about 1960, although the number of cold days has continued to fall. Given the similar timing of the increase in numbers of warm days (Figure 1.2a), it is reasonable to assume that the decline in numbers of cold days over recent decades has the same causal mechanism.

2003 was the 4th hottest summer in the CET record. It was exceptional in terms of absolute maxima (with new record maxima for both England and Scotland) and for the extent and persistence of very high night-time temperatures (Burt, 2004a,b). On 10th August 2003, maximum temperatures greater than 37°C were recorded at at least 13 locations in Southeast England (Burt, 2004a). Analyses of the very hot summer of 2003 from a climatic perspective have, however, tended to focus on France, Germany, Italy and, in particular, Switzerland (Beniston, 2004; Luterbacher *et al*, 2004; Schar *et al*, 2004; Stott *et al*, 2004). These studies indicate that the summer of 2003 was probably

the hottest in Europe since the year 1500 and that human influence has at least doubled the risk of a heatwave of this magnitude occurring.

Figure 1.2a Number of days per year with daily mean CET exceeding 20° C





Figure 1.2b Number of days per year with daily mean CET below 0° C



1.2.3 Rainfall

There has been a rising trend in annual total rainfall in England and Wales since about 1970, but this is well within the natural variability found over the series as a whole (Wigley *et al*, 1984; Jones and Conway, 1997).

When rainfall is broken down into its seasonal components, however, pronounced linear trends are identified in all seasons apart from spring. Summer rainfall has fallen by about 50 mm over the last 250 years, whereas winter and autumn rainfall have increased by about the same amount over this period. As with cold temperatures (Section 1.2.2), these trends are consistent over much too long a period to be associated fully with global warming. It is possible that for the first part of the record the trend is due to recovery from the “Little Ice Age”. Given the absence of abrupt changes in rainfall regime over the last few decades, such as those described for temperature, it is not possible to attribute any of the recent England and Wales precipitation trend to anthropogenic warming.

The England and Wales rainfall series discussed above allows examination of long-term rainfall variability, but does not tell us anything about spatial variability in rainfall trends. Analysis of 146 UK weather-station records for the period 1961–2000 indicates increased rainfall totals and increased frequency and contribution of heavy rainfall events during winter, and decreases during summer (Osborn and Hulme, 2002). These changes tend to be spatially coherent, i.e. broadly similar across the UK, although the largest winter changes tend to occur in the Western UK (especially Northwest England and Western Scotland) and summer rainfall decreases are less evident in Western Scotland, South Wales and parts of Northern Ireland. [Changes in rainfall due to climate change are discussed in relation to flood risk in Chapter 3].

1.3 Future climate change in the UK: the UKCIP02 scenarios

The world's changing climate is a result of both natural and anthropogenic factors. A stable climate requires a balance between incoming solar radiation and outgoing radiation. Fundamental to this stable state is the natural greenhouse effect, the process whereby greenhouse gases such as carbon dioxide, methane and nitrous oxide absorb and emit infrared radiation, trapping heat within the atmosphere and maintaining a suitable temperature for human habitation.

Anthropogenic climate change is caused by the rising concentration of greenhouse gases and aerosols in the atmosphere. These arise from various human activities performed with increasing intensity since the beginning of the Industrial Revolution in the late 18th Century. Natural variations in climate have historically been associated with a number of factors, including: variations in solar output, changes in the orbital characteristics of the Earth, and volcanic eruptions - ash and sulphur dioxide emissions alter the reception and reflection patterns of solar radiation. However, the balance of evidence indicates that that warming experienced in the last 50 years is "very likely" to be due to human activities (Houghton *et al*, 2001).

The UK is fortunate in having national climate scenarios (Hulme and Jenkins, 1998; Hulme *et al*, 2002). The current set of scenarios – called UKCIP02 – are based upon the latest socioeconomic scenarios (the SRES, see Nakicenovic *et al*, 2000) and were constructed from a high-resolution Regional Climate Model (RCM) HadRM3h, developed by the Hadley Centre¹. For these and other reasons, the UKCIP02 scenarios represent a significant improvement in scenario construction. Box 1.1 summarises the main differences between the UKCIP02 and the UKCIP98 scenarios which were used in the first climate change and health report (Department of Health, 2001).

It is important to be aware that climate models such as those used in the construction of the UKCIP scenarios do not provide a definitive prediction of what climate will be like at a particular time in the future. The models simulate climate change according to a particular set of assumptions about the world, including a particular regimen of greenhouse gas emissions.

There are two main categories of uncertainties in all climate scenarios: future emissions of greenhouse gases which depend on future human activities; and scientific uncertainties. The latter are, in theory, reducible. Climate model uncertainties, for example, can, in part, be addressed by considering output from a number of different models rather than from a single model. The UKCIP02 scenarios are based on output from a single suite of models, although some intercomparisons with other models are presented in the Scientific Report (Hulme *et al*, 2002) and the Hadley Centre has subsequently undertaken some more detailed intercomparisons with output from other RCMs (Jenkins and Lowe, 2003; Rowell, 2004). The most recent Hadley Centre commentary on the UKCIP02 scenarios considers the effect of global dimming² (i.e. a reduction in global radiation at the surface due to changes in atmospheric aerosols and clouds). While the underlying HadCM3 simulations do show global dimming over the past few decades, this is only a small fraction of that observed.

The UKCIP02 scenarios are described in a scientific report (Hulme *et al*, 2001) and summarised in a briefing report (both available from the UKCIP website). These reports contain many maps showing the scenario changes. Most of these show a North-West to

¹ The Hadley Centre for Climate Prediction and Research, the Met Office, UK.

² http://www.ukcip.org.uk/news_releases/28.pdf

South-East gradient, with the largest changes occurring in the Southeast. Table 1 summarises the changes in mean winter and summer temperature and rainfall for this region of largest change.

Table 1.1 UKCIP02 scenario changes in mean seasonal temperature and rainfall for Southeast England (taken from Figures 31-38 in Hulme *et al*, 2002)².

	2020s	2020s	2050s	2050s	2080s	2080s
	Winter	Summer	Winter	Summer	Winter	Summer
Temperature °C						
Low	0.5 to 1	1 to 1.5	1 to 1.5	2 to 2.5	1.5 to 2	2.5 to 3
Medium-low	0.5 to 1	1 to 1.5	1 to 1.5	2 to 2.5	1.5 to 2	3 to 3.5
Medium-high	0.5 to 1	1 to 1.5	1.5 to 2	2.5 to 3	2.5 to 3	>4.5
High	0.5 to 1	1 to 1.5	1.5 to 2	3 to 3.5	3 to 3.5	>4.5
Rainfall %						
Low	0 to +10	-10 to -20	+10 to +15	-10 to -30	+15 to +20	-20 to -30
Medium-low	0 to +10	-10 to -20	+10 to +15	-20 to -30	+15 to +20	-30 to -40
Medium-high	0 to +10	-10 to -20	+10 to +15	-20 to -30	+20 to +30	-40 to -50
High	0 to +10	-10 to -20	+15 to +20	-30 to -40	+25 to >+30	> -50

Box 1.1. UKCIP02 vs. UKCIP98

The differences in the future climates between UKCIP98 and UKCIP02 are due to the use of different climate models, and different assumptions about future greenhouse gas emissions, particularly the upper estimate of “high” greenhouse gas emissions.

- UKCIP02 scenarios are slightly warmer than UKCIP98 because they are based on generally higher carbon dioxide emissions and lower sulphur dioxide emissions (the latter cause cooling rather than warming), and the HadCM3 model tends to be more sensitive to the greenhouse gas forcing (see Table 3 in Hulme *et al*, 2002).
- However, changes in the representation of thermal expansion of sea water and the dynamics of glacier melting, mean that the UKCIP02 scenarios show slightly less sea-level rise than UKCIP98.
- The UKCIP02 scenarios suggest that summers may become drier across the whole of the UK, not just in England and Wales, and by a larger amount than UKCIP98. UKCIP02 also indicates slightly drier conditions in spring and autumn, rather than wetter as UKCIP98. For Scotland, the UKCIP02 scenarios show significantly different patterns in rainfall change, with a much smaller increase in autumn rainfall than UKCIP98.
- There are also some differences in the pattern of changes in wind between the UKCIP98 and UKCIP02 scenarios, but despite the model improvements, the UKCIP02 changes in wind are considered so uncertain, that no level of confidence can be assigned to them (whereas changes in other variables are ranked in terms of their relative confidence: high, medium and low).
- The use of a suite of models providing a range of integrations allows a more rigorous analysis of uncertainties and extreme events.

² See Table 5 in Hulme *et al*, 2002 for uncertainty margins at the 50 km grid-box scale.

1.4 Future climate change in the UK: extreme events

The UK is projected to experience substantial changes in extreme climate events due to anthropogenic climate change by the end of this century:

- warmer winters and hotter summers
- widespread increases in the duration of hot spells in parts of England
- extended summer drought periods over much of England and Wales
- lower extreme wind speeds

The recent availability of output from a number of European RCMs has allowed new analyses focusing on climate extremes which are important to human health (heavy rainfall, hot and cold spells). Results are discussed for the UK using the available RCM data³ for two time periods: 1961-90 (current climate) and 2070-99. All results presented here are for one emissions scenario – the A2 scenario (Nakicenovic *et al*, 2002) which is comparable to the UKCIP02 medium-high scenario.

The analyses of future UK climate use data from three RCMs:

- HadRM3P – Hadley Centre, UK Met Office (Hadley Centre, 2002 and Hudson and Jones, 2002)
- HIRHAM4 – Danish Meteorological Institute (Christensen *et al*, 1996)
- RCAO – Rossby Centre, Swedish Meteorological and Hydrological Institute (Döscher *et al*, 2002).

Validation of the RCMs is described elsewhere (e.g. Holt *et al*, 2005 for HadRM3P). In general, annual indices of extremes compare reasonably well between NCEP Reanalysis data and HadRM3P, although the model extremes of temperature and wind speed generally compare better than rainfall. Duration extremes, such as longest spell of drought, compare better than indices based on absolute values.

All three RCMs were forced with output from the HadAM3P high-resolution GCM (a slightly updated version of the GCM used to construct the UKCIP02 scenarios). At the start of the modelling chain, HadAM3P was driven by output from the HadCM3 coupled atmosphere-ocean GCM. For the A2 emissions scenario, HadCM3 gives a global temperature change of 3.3°C by 2100. This is in the middle of the 1.4-5.8° C range quoted in the last IPCC assessment report (Houghton *et al*, 2001) based on the full range of 35 SRES scenarios and a number of climate models.

We consider temperature, rainfall, and wind speed extremes represented by a single maximum value per year. The change in climate is assessed by comparing the change in mean value of the climate extreme between the periods 1961-90 and 2070-99, averaged over the ensemble of three models. The cross-model average is known as the ensemble mean. Uncertainty in the ensemble mean is assessed using a bootstrap method in which the data from all models are re-sampled with replacement 1000 times and the difference between means calculated on each iteration. The confidence limits at the 5% significance level are then defined by the 25th and 975th values of the sorted 1000 differences (Davison and Hinkley, 1997).

The indices fit the Generalised Extreme Value (GEV) distribution (Coles, 2001) which enables the calculation of robust estimates of return levels for different return periods. We compare the difference in return levels between the present and the 2080s. For example, return levels of annual maximum length of dry spell for a 100 year return

³ This problem was overcome in the UKCIP02 scenarios by using a technique called pattern-scaling. However, this approach is not appropriate for application to the climate extremes considered here.

period, calculated from a GEV distribution using 1961-90 data, have a 1% chance of occurring in a given year. Given the probability that warming will lead to extended dry periods in 2070-99, the 100-year event calculated using data from the end of the century will be considerably longer than at present. Similarly, the current 100-year event will have a much higher chance of occurring at the end of the century than 1% in a given year.

Projected changes in temperature, rainfall and wind extremes are presented in Sections 1.4.1.1 to 1.4.4. It is stressed that these are based on a single emissions scenario (A2/medium-high). They are also based on RCM simulations forced by one underlying set of GCM simulations. It has been demonstrated that inter-model variability associated with the choice of driving models is a major source of the scientific uncertainty (Christensen *et al*, 2006). In Section 1.5, we consider how the probability of getting heatwaves of the order of the 2003 event might evolve over the coming century.

1.4.1 Temperature extremes

A standard format is used to illustrate changes in climate. A map of the projected climate change (Figure 1.3a) is accompanied by a map of the uncertainty in the change (Figure 1.3b). The uncertainty is determined at the 5% significance level.

Changes in annual mean temperature. The projected change in mean annual temperature over most of the UK (Figure 1.3a) is generally less than 3° C, with the greatest increases being found in the South. The uncertainty in these estimates (Figure 1.3b) is quite low, being typically $\pm 5\%$ of the projected change in temperature for a particular area. Looking at Southeast England in Figure 1.3a, we can see that an area roughly covering Kent, Surrey, and Hampshire could expect mean annual temperatures to rise by between 3 and 4.5° C under the A2 scenario by the 2080s. Looking at the uncertainty in Figure 1.3b, we find this region to have uncertainty of ± 0.1 to 0.15° C. Therefore, concentrating for the sake of simplicity on the upper end of the range, we can say that mean annual temperature in this area is projected to increase by $3.5^\circ \text{ C} \pm 0.15^\circ \text{ C}$, or that 95% of the time mean annual temperatures at the end of the century would be between 3.35 and 3.65° C warmer than today, under the A2 scenario.

The changes in mean summer temperatures are considerably larger than those for winter. The biggest summer increases of between 3.5 and 4° C ($\pm 0.3^\circ \text{ C}$) are projected for virtually the whole of Southern England, while winter changes are generally less than 2.2° C .

Figure 1.3a Change in mean annual temperature (°C) between 1961-90 and 2070-99, under the A2 scenario (darker red is warmer in future).





Figure 1.3b Uncertainty ($\pm$ the amount shown in °C) in the change in mean annual temperature shown in Figure 1.3a (pink is greater uncertainty).



Change in duration of hot and cold spells

Cold spells. Figure 1.4a shows the projected change in the longest spell in a year with daily minimum temperature below freezing. This cold spell is expected to become shorter over the whole of the UK. The biggest changes are to be found in Scotland where the main freezing period is projected to shorten by about a week. The small changes over Ireland and the coast of the UK are due to these maritime areas generally not having freezing episodes over long periods. The uncertainty in the projected change in the length of cold spell is quite high (Figure 1.4b) at about $\pm 16\%$.

Figure 1.4a Change in longest spell (days) with daily minimum temperature below 0°C between 1961-90 and 2070-99, under the A2 scenario (darker blue is least shortening of cold spells in the future).






Figure 1.4b Uncertainty ($\pm$ the amount shown in days) in the change in the longest length of cold spell shown in Figure 1.4a (pink is greater uncertainty)



Hot spells. Projected changes in the length of hot spells, defined as the longest spell with temperatures above 25° C are shown in Figure 1.5a. Note that this climatic definition of a hot spell does not correspond to an actual heatwave as might be defined in health impact terms which would be a more extreme event of shorter duration. Generally, only Central and Southern England experience periods with grid-box temperatures as high as this, but these hot spells are projected to increase by between 5 to 10 days in places. The uncertainty in these estimates is about ± 10 -14% (Figure 1.5b).

Note in Figure 1.5a that large changes in the duration of high temperature episodes are indicated for Northern France. Studies over the whole of France indicate that extreme heatwaves similar to the 2003 episode are likely to become common over Central and Southern France.

Figure 1.5a Change in longest spell (days) with daily maximum temperature above 25°C between 1961-90 and 2070-99, under the A2 scenario (darker red is longer warm spells in the future). **Figure 1.5b Uncertainty ($\pm$ the amount shown in days) in the change in the longest length of hot spell shown in Figure 1.5a (pink is greater uncertainty)**



1.4.2 Rainfall extremes

Rainfall is one of the most difficult physical parameters to model. This is reflected in the relatively large uncertainty in the analyses below.

The projected changes in annual total rainfall shown in Figure 1.6a give a mixed picture for the UK. Generally, the wetter North and Ireland are projected to become drier and the Southern coast is likely to become wetter. However, even the biggest of these changes is quite small, being typically less than 7% of the rainfall total in a particular region. Given the very high uncertainty in these projections (Figure 1.6b), often $\pm 60\%$ of the projected change, it is difficult to assess their severity. For example, applying the upper bound uncertainty assessment, Scotland and Ireland can expect to lose about 12% of their annual total rainfall by the end of the century.

Figure 1.6a Change in total annual rainfall (mm) between 1961-90 and 2070-99, under the A2 scenario (darker green is wetter in future).





Figure 1.6b Uncertainty ($\pm$ the amount shown in mm) in the change in total annual rainfall shown in Figure 1.6a (pink is greater uncertainty).



Change in length of longest wet and dry spells

The projected changes in the maximum length of dry spell are shown in Figure 1.7a. The worst affected areas are Southern and Western England with an extension of the longest period of drought by between 4 and 6 days. Taking account of the large uncertainty (Figure 1.7b) could extend this by 2 days. In itself, such an increase is unlikely to stress water supplies unduly. However, since this is an average, it indicates that the most extreme droughts could be considerably longer than at present. This is considered further in the examination of changes in return levels in Section 4.1.4.

Figure 1.7a Change in maximum length of dry spell (days) between 1961-90 and 2070-99, under the A2 scenario (red is longer dry spells in the future).




Figure 1.7b Uncertainty ($\pm$ the amount in days) in the change in the maximum length of dry spell shown in Figure 1.7a (pink is greater uncertainty).





Figure 1.8a shows projected changes in the longest period of rainfall in a year. Apart from a few coastal areas, this is shown to shorten by a couple of days over most of the country. The uncertainty in this estimate is very high, reaching $\pm 70\%$ in places (Figure 1.8b).

Figure 1.8a Change in maximum length of wet spell (days) between 1961-90 and 2070-99, under the A2 scenario (blue is longer wet spells in the future).

Figure 1.8b Uncertainty ($\pm$ the amount shown in days) in the change in the maximum length of wet spell shown in Figure 1.8a (pink is greater uncertainty)




1.4.3 Maximum daily wind speed

The direct impact of wind on human health is caused by high wind speeds (see Chapter 2). Maximum daily wind speed is the highest wind speed in a day occurring instantaneously on a model timestep, and has no equivalent in observed meteorological data. For this reason, a number of RCMs, although none of those considered here, are starting to include an estimate of maximum gust, intended to simulate the 10 minute gust speed recorded by anemometers.

The projected change in the annual highest maximum daily wind speed (Figure 1.9a) is relatively trivial over most of the UK, particularly considering the relatively large uncertainty (Figure 1.9b). Most of the country is projected to experience slightly higher wind speeds in the most severe storms. The greatest increases in wind speeds are indicated over Southeast England, Central Ireland, and Northern Scotland.

Figure 1.9a Change in annual highest maximum daily wind speed (m/s) between 1961-90 and 2070-99, under the A2 scenario (darker brown is higher future wind speeds).

Figure 1.9b Uncertainty ($\pm$ the amount shown in m/s) in the change in the annual highest maximum wind speed shown in Figure 1.9a (pink is greater uncertainty)



1.4.4 Projected changes in return levels

Calculation of return levels for periods longer than the length of record necessarily involves extrapolation. This is supported by extensive theory, based on the fit of the data to the GEV distribution, and the validity of certain underlying assumptions about the statistical properties of the data (Coles, 2001). Beyond certain somewhat arbitrary “rules of thumb”, there appears to be no robust definition of what is the maximum reliable return period for a given length of record. Here we use 100 years because it is convenient, commonly used, and only slightly over three times the length of the record.

The calculation of return levels is successful only if the data on each RCM grid square fit the GEV distribution, defined by mean, variance, and shape parameters (Coles, 2001). The three models used here do not always yield valid GEV parameters at the same grid points for each extreme index. Generally, the HadRM3P model provides more consistent GEV parameters than the HIRHAM4 model which, in turn, is better in this regard than the RCAO model. The discrepancies between the models make it impossible to examine a valid ensemble mean return level. Therefore, we present return levels for the HadRM3P model only, bearing in mind that the results have none of the safeguards of an intermodel comparison and that it is not possible to provide a meaningful estimate of uncertainty.

Cold spells


Figure 1.10a shows the 100-year return level for cold spells over the UK using 1961-90 data. A one in a 100-year cold spell is approximately 20-30 days with temperatures less than 0° C in Southern England for the baseline climate. Cold spells lasting between 60 and 70 days can be expected, on average, once every 100 years over Central and Western Scotland. Under the A2 scenario of climate change, the length of 100-year return period cold spells in Scotland is projected to be reduced by up to 40 days (Figure 1.10b).

Just as a return level can be calculated from the parameters of the GEV distribution and a specified return period, so can a return period be calculated from the GEV and a specified return level. This gives us a slightly different way of looking at Figures 1.10a and 1.10b which may be considered more informative.

Using the midpoint of each data “bin”, according to the 1961-90 data (Figure 1.10a) there is a 1% chance of having a cold spell of at least 65 days in Scotland in a given year. Under climate change, however, the 2070-99 data (Figure 1.10b) shows a 1% probability of a cold spell lasting at least 35 days in Scotland in a given year. This dramatic reduction in the risk associated with a low probability event by the end of the century can also be considered in terms of today’s climate. A 35-day cold spell in Scotland would currently be quite common and expected about once every five years, or have a 20% chance of occurring in a given year (compared with 1% by the end of the century).

Figure 1.10a 100-year return level (days) for the annual longest period with daily minimum temperatures less than 0° C. Based on 1961-90 extremes (pale blue is the longest cold period). Using HadRM3P data only.

Figure 1.10b Projected change, under the A2 scenario, in the 100-year return level (days) for the annual longest period with daily minimum temperatures less than 0°C (pink shows the greatest warming).




Prolonged hot spells

Figures 1.11a and 1.11b present changes in return levels of the longest annual hot spell (defined as a period with daily maximum temperatures above 25°C for each 90km grid square), according to 1961-90 and 2070-99 data, respectively.

There is a 1% chance of a hot spell longer than 35 days in a given year over Central Southern England, according to 1961-90 data (Figure 1.11a). By the 2080s, this area has a 1% chance of hot spells longer than 60 days in a given year (Figure 1.11b). Currently, there is only a 0.01% chance of hot spells exceeding 60 days in Southern England in a given year.

Figure 1.11a 100-year return level (days) for the annual longest period with daily maximum temperature above 25° C. Based on 1961-90 extremes (pink is the longest warm period). Using HadRM3P data only.

Figure 1.11b Projected change, under the A2 scenario, in the 100-year return level (days) for the annual longest period with daily maximum temperatures above 25° C (pink shows the greatest warming).




Wet and dry spells

The return levels for prolonged wet spells, based on 1961-90 data (Figure 1.12a), show that apart from a few western extremities with a 1% chance of wet spells in excess of 85 days, most of the country has a 1% chance of spells of around 25 days. The effects of climate change on the duration of severe wet spells is complex. Using data from the 2070-99 period (Figure 1.12b), most of the country sees a decline in the length of 1% wet spells by about 30%. The islands off the west of Scotland, however, see 1% wet spells increase in length to about 100 days by the end of the century. This is equivalent to a 1 in 400 year event now, or a chance of about 0.025% of wet spells occurring in a given year.


Figure 1.12a 100-year return level (days) for the annual longest period with rainfall every day. Based on 1961-90 extremes (blue is the longest wet period). Using HadRM3P data only.

Figure 1.12b Projected change, under the A2 scenario, in the 100-year return level (days) for the annual longest wet spell (pale blue shows longer wet period).



The 100-year drought (defined as the maximum number of consecutive days without rain) is currently about 45 days for the UK (Figure 1.13a). The average increased length of drought is about 10 days under global warming (Figure 1.13b), apart from Central Southern England, which is consistent with expectations of longer hot spells. Here, there is a 1% chance of drought exceeding 75 days in a given year by the end of the century. This is equivalent to a 0.013% chance of drought occurring under current climate conditions. Given these projections, water shortages are likely to become a regular feature of UK summers unless infrastructure changes are planned now.

Figure 1.13a 100-year return level (days) for the annual longest period with no rainfall based on 1961-90 extremes (yellow is the longest dry extremes). **Figure 1.13b Projected change, under the A2 scenario, in the 100-year return level (days) for the annual longest dry spell (pink shows biggest extension of dry period).** Using HadRM3P data only.



Highest maximum wind speed

The highest 100-year return levels for annual highest daily maximum wind speed based on 1961-90 data are around the coast of the UK (Figure 1.14a), with isolated locations exceeding 40 m/s. The effects of climate change (Figure 1.14b) are quite complex with most parts of the UK experiencing lower extreme wind speeds by the end of the century, apart from increases over much of Wales, Northeast England and Northwest Scotland. Over much of Northern Ireland, for example, current 1% maximum wind speeds of 25 m/s decline to 15 m/s by the end of the century. This is equivalent to a 10% chance of occurrence in a given year under the climate of today, a much less hazardous event. Over Wales, future 1% maximum winds are about 5 m/s stronger than at present, equivalent to 0.025% winds under current climate.

Figure 1.14a 100-year return level (m/s) for the annual highest daily maximum wind speed. Based on 1961-90 extremes (darker brown is the highest wind speed). Using HadRM3P data only.





Figure 1.14b Projected change, under the A2 scenario, in the 100-year return level (m/s) for the annual highest daily maximum wind speed (pink shows higher future wind speed pale blue is lower future wind speeds).



1.5 Changes in the frequency of heatwaves of the order of the 2003 event

Following the events of 2003, the UK has introduced a heatwave plan which includes a system for triggering alerts as soon as the Met Office forecasts ‘threshold temperatures’ for at least three days ahead in one region, or that there is an 80% chance of temperatures being high enough on at least two consecutive days to have significant effects on health. For the London region, the operational day-time threshold temperature, is 32° C. It would clearly be desirable to have projections of how the frequency of occurrence of these threshold temperatures will change for each region. The grid-box nature of climate model output means, however, that thresholds defined from point locations are not directly comparable to the model grid boxes. Thus further downscaling would be required in order to construct such scenarios. Here, however, we explore future changes in the frequencies of heatwaves of the order of the 2003 episode using a simple yardstick applied to output from a GCM simulation for the period 1990-2100.

GCM output is used for this analysis because it allows us to examine the evolution over time of the impacts of climate change. RCMs offer undeniable improvements in physics and reliability over the coarser resolution GCMs. They have a much higher grid resolution – typically 50 km, rather than the 300 km typical of GCMs. However, currently output is only available for ‘snapshot’ periods, i.e. 1961-90 and 2070-99 (see Section 1.4).


We use data from the UK Met Office HadCM3 global model from 1990 to 2100 to provide estimates of changes in the frequency of heatwaves of the order of the 2003 event under the A2 emissions scenario. The model data are interpolated onto a 2 x 2 degree latitude/longitude grid to give improved coverage over the UK. A point roughly located at Cambridge is selected for the analysis. It is not possible to select a point further south

because of the coarse grid. However, the climate of Cambridge is broadly representative of the Southeast of England. To give a yardstick against which to compare the model estimates, we select data for the Cambridge location from the NCEP reanalysis for 2003. The reanalysis is based on observational data from a variety of sources and is placed on a grid of roughly 2.2 degrees latitude/longitude. The gridding process smoothes the data, making it more suitable for comparison with the gridded model data than point observations from stations.

The variable we chose to analyse is a 7-day running mean daily maximum temperature at the surface. This is a sufficiently long averaging period to indicate a persistent hot spell. The NCEP reanalysis yielded six days when temperatures averaged over 7 days were 25°C in 2003, the hottest running mean in the year for Cambridge. Comparison of NCEP maximum temperature percentiles over Cambridge with HadCM3 percentiles, suggest that the model data are about 2°C warmer than NCEP. Thus, to compare HadCM3 with NCEP, we impose a penalty of 2°C and search for days when the 7-day running mean from HadCM3 was 27°C or higher.

The results of the analysis are shown in Figure 1.15. The six days with a 7-day running mean of at least 25°C from the NCEP data are shown by the pink continuous line at the bottom of the plot. This is the yardstick against which the severity of future warming can be assessed. Bearing in mind that 27°C from the model running mean is equivalent to the NCEP 25°C, it is clear that the equivalent of the 2003 event for the South of England will be encountered occasionally until about 2030 (approximately 4 times in the 40 year period). The first three decades of the century are shown to be no different from the present in terms of variability. After this, however, heatwaves such as that of 2003 can be expected much more frequently, becoming increasingly more severe in intensity and duration after 2060.

Figure 1.15 Number of days per year when the 7-day running mean daily maximum temperature from HadCM3 is at least 27°C. The pink line shows the comparable six days above 25°C from the NCEP data that was achieved in the 2003 heatwave.



1.6 Low probability, high impact events

Low probability, high impact events are, by definition, rare. Such events can have a number of different climatic causes and impacts. Certain geophysical events are also of concern as these may impact indirectly on the climate of the British Isles. The major geophysical events of concern are large volcanic eruptions, sometimes referred to as super volcanoes. Such eruptions, especially those that occur in the tropics, can have a cooling impact upon the global climate. Ash and debris emitted into the atmosphere can cause a veil of reflective particles that reduces the amount of incoming short-wave radiation from the sun, thus causing cooling. It is not, however, possible to predict whether such a volcanic event will occur in the coming century.

The major climate events that we are most concerned with in terms of impacts on the British Isles are: a shutting down of the North Atlantic thermohaline circulation; a collapse of the West Antarctic Ice Sheet; and accelerated climate change as a result of the release of naturally stored greenhouse gases.

The first of these, thermohaline circulation collapse, has been very much discussed and was popularised in the Hollywood film *'The Day after Tomorrow'*. Palaeoenvironmental data suggests that there has been a number of abrupt changes in the climate of the Northern Hemisphere during the last 250,000 years. These changes are believed to have occurred as a result of the input of a large amount of freshwater into the North Atlantic, caused by rapid melting of a large (or series of) ice sheets. The input of freshwater breaks down the thermohaline circulation (THC), thus deflecting (or stopping) the North Atlantic Drift (Gulf Stream). The Gulf Stream acts to warm North West Europe, providing it with a mild maritime climate which is some 9°C warmer than the average for this latitude. Thus, during periods of THC shutdown, the absence of the Gulf Stream results in a colder continental-type climate for North West Europe, including the British Isles.

It has been hypothesised that during a period of warming (such as that which we are experiencing now and is projected for the future) there may be a reoccurrence of the shutdown of the THC as a result of increased freshwater inputs due to the melting (or partial melting) of the Greenland Ice Sheet and an enhanced hydrological cycle. Observations are showing a slight weakening in the strength of the THC, while experiments using coupled ocean-atmosphere GCMs have shown that the current generation of climate models are able to induce a shutdown of the THC and associated cooling if large amounts of freshwater are artificially added to the North Atlantic. To date, however, no GCM experiment has produced a shutdown of the THC when forced with realistic scenarios of greenhouse-gas-induced climate change, although most GCMs, including the Hadley Centre simulations used in Sections 3 and 4, project a general weakening of the THC. It must also be taken into account that if the THC were to shutdown, it would be against the backdrop of very much higher greenhouse-gas-induced climate change. The scientific consensus is that a shutdown of the THC is unlikely without the rapid collapse of the Greenland Ice Sheet and that this is highly unlikely to occur before 2100 within the range of temperatures projected by the current generation of SRES forced GCM experiments.

Through the thermal expansion of sea water and the melting of small (alpine type) glaciers, global sea-level rise has been occurring in line with human induced warming. Future rates of change under the SRES scenarios suggest that during the 21st century sea-level rise will be in the order of 20-80 cm (Houghton *et al*, 2001). There is concern, however, that more rapid rates of sea-level rise are likely as a result of the collapse of the West Antarctic Ice Sheet (WAIS). Evidence from satellite sensor data has shown areas of weakness in the WAIS which some scientists have suggested could indicate that it is relatively unstable and could be prone to collapse. If such a collapse occurs, then global sea level would rise by some 8 metres over the course of a number of centuries. At present, there is no consensus on the timing or possibility of such rapid sea-level rise, though it is generally considered highly unlikely to occur in the present century.

Climate change as a result of human activities is causing rapid rates of temperature rise and there is some concern that this may trigger positive non-linear feedbacks in the climate system. Such feedbacks could affect both the biogeochemical and physical systems. The natural environment stores large amounts of greenhouse gases in reservoirs, primarily in the form of methane and carbon dioxide. It is a concern, therefore, that climate change may cause the release of these gases and thus enhance their concentrations in the atmosphere. The major reservoirs can be found: locked in organic matter in the permafrost of the Northern Hemisphere; in the terrestrial carbon sink e.g. the Amazonian rain forest; and, locked in particulate matter on the ocean floor. Recent monitoring has shown that the permafrost regions are starting to melt and are releasing carbon dioxide and methane. Modeling work undertaken by the Hadley Centre has shown that, as a result of increased temperatures, the Amazon is likely to become a source of carbon dioxide by the end of the century due to forest die back. Rapid release of methane from deep ocean deposits is also a feasible scenario. However, as yet, there is little evidence to suggest when and if this is likely to happen.

The major physical feedback that is likely to accelerate climate change which is factored into climate models, is reduction in snow and ice cover in the high latitudes. Snow and ice have a high albedo and thus reflect much of the incoming solar radiation. As a result of warming, sea-ice extent (especially in the Arctic) is reducing, thus lowering the albedo and causing more of the incoming solar energy to be absorbed at the surface. This in turn leads to higher temperatures, increased reduction of sea-ice and more warming i.e. a positive feedback effect. The Arctic sea-ice extent during the summer of 2005 was at its minimum level yet recorded.

Acknowledgements

This research would not have been possible without the data made freely available by the following institutions:

- The climate model data used in this study were downloaded from the PRUDENCE Web Site at the Danish Meteorological Institute: <http://prudence.dmi.dk/>
- The Central England Temperature Series and the England & Wales Rainfall data are available from the British Atmospheric Data Centre: <http://badc.nerc.ac.uk/home/> and from the Hadley Centre: <http://www.metoffice.com/research/hadleycentre/>
- UKCIP02 scenario data are available from the UKCIP02 Scenario Gateway at <http://www.ukcip.org.uk/resources/tools/scenarios.asp>

References

- Beniston, M. (2004) The 2003 heat wave in Europe: A shape of things to come? *Geophysical Research Letters* **31**, L02202, doi:10.1029/2003GL018857.
- Burt, S. (2004a) The August 2003 heatwave in the United Kingdom: Part 1 – Maximum temperatures and historical precedents. *Weather* **59**, 199-208.
- Burt, S., 2004b: The August 2003 heatwave in the United Kingdom: Part 3 – Minimum temperatures. *Weather* **59**, 272-273.
- Christensen, J. H., Christensen, O.B., Lopez, P., Van Meijgaard, E. and Botzet, M. The HIRHAM4 Regional Atmospheric Climate Model, *Scientific Report 96-4*, 51 pp., DMI, Copenhagen, 1996.
- Christensen, J.H. *et al*, (2006) Evaluating the performance and utility of regional climate models in climate change research: Reducing uncertainties in climate change projections – the PRUDENCE approach, *Climatic Change*, in press.
- Coles, S. (2001) *An Introduction to Statistical Modeling of Extreme Values*. Springer Series in Statistics, Springer-Verlag, London, ISBN 1-85233-459-2, 208 pp.
- Davison, A.C. and Hinkley, D.V. (1997) *Bootstrap Methods and Their Applications*. Cambridge Series in Statistical & Probabilistic Mathematics, Cambridge University Press; ISBN: 0521574714, 592 pp.
- Döscher, R., Willén, U., Jones, C., Rutgersson, A., Meier, H. E. M., Hansson, U. and Graham, L. P. (2002) The development of the coupled regional ocean-atmosphere model RCAO. *Boreal Env.Res.* **7**, 183-192.
- Hadley Centre. (2002) *The Hadley Centre regional climate modelling system: PRECIS update 2002*. Downloadable web document from the UK Meteorological Office: <http://www.metoffice.com/research/hadleycentre/pubs/brochures/B2002/precis.pdf>.
- Holt, T., Barring, L., Kundzewicz, Z.W. and Palutikof, J.P. (2005) Comparison of extremes from global and regional climate models with NCEP reanalysis data. Submitted to *Climate Dynamics*.
- Houghton, J.T., Ding, Y., Griggs, D.J., Noguer, M., van der Linden, P.J., Dai, X., Maskell, K. and Johnson, C.A. (eds.), (2001) *Climate Change 2001: The Scientific Basis*, Cambridge University Press.
- Hudson, D.A. and Jones, R.G. (2002) *Regional Climate Model Simulations of Present-Day and Future Climates of Southern Africa*. Hadley Centre Technical Note 39, Hadley Centre, Met Office, Exeter, UK, 41 pp.
- Hulme, M. and Jenkins, G.J. (1998) *Climate change scenarios for the United Kingdom*. UKCIP Technical Note No. 1., Climatic Research Unit, Norwich.
- Hulme, M., Jenkins, G.J., Lu, X., Turnpenny, J.R., Mitchell, T.D., Jones, R.G., Lowe, J., Murphy, J.M., Hassell, D., Boorman, P., McDonald, R. and Hill, S. (2002) *Climate Change Scenarios for the United Kingdom: The UKCIP02 Scientific Report*, Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia, Norwich, UK, 120pp.

- Jenkins, G. and Lowe, J. (2003) *Handling uncertainties in the UKCIP02 scenarios of climate change*, Hadley Centre technical note 44: http://www.met-office.gov.uk/research/hadleycentre/pubs/HCTN/HCTN_44.pdf
- Jones, P.D. and Conway, D. (1997) *Precipitation in the British Isles: an analysis of area-average data updated to 1995*. International Journal of Climatology, **11**, 331-345.
- Luterbacher, J., Dietrich, D., Xoplaki, E., Grosjean, M. and Wanner, H. (2004) European seasonal and annual temperature variability, trends, and extremes since 1500. *Science* **33**, 1499-1503.
- Manley, G. (1953) *The mean temperature of Central England, 1698 to 1952*. Quart J Roy Meteorol.Soc. **79**, 242-261.
- Manley, G. (1974) *Central England Temperatures: monthly means 1659 to 1973*. Quart J Roy.Meteorol.Soc. **100**, 389-405.
- Nakicenovic, N., Alcamo, J., Davis, G. et al. (2000) *"Emissions Scenarios : A Special Report of Working Group III of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge.
- Osborn, T.J. and Hulme, M. (2002) Evidence for trends in heavy rainfall events over the UK. *Philosophical Transactions of the Royal Society of London A*, **360**, 1313-1325.
- Parker, D.E., Legg, T.P. and Folland, C.K. (1992) A new daily Central England Temperature Series, 1772-1991. Int. J. Climatol. **12**, 317-342.
- Rowell, D.P. (2004) *An initial estimate of the uncertainty in UK predicted climate change resulting from RCM formulation*, Hadley Centre technical note 49: http://www.met-office.gov.uk/research/hadleycentre/pubs/HCTN/HCTN_49.pdf
- Schär, C., Vidale, P.L., Lüthi, D., Frei, C., Häberli, C., Liniger, M.A. and Appenzeller, C. (2004) The role of increasing temperature variability in European summer heatwaves. *Nature* **427**, 332-336.
- Stott, P.A., Stone, D.A. and Allen, M.R. (2004) Human contribution to the European heatwave of 2003. *Nature* **432**, 610-614.
- Wigley, T.M.L., Lough, J.M. and Jones, P.D. (1984) Spatial patterns of precipitation in England and Wales and a revised homogeneous England and Wales precipitation series. *Journal of Climatology* **4**, 1-25.

2 Flooding and windstorms and climate change

(See also section 4.5. – windstorms & section 4.6 – flooding of 2001 report)

Sari Kovats, London School of Hygiene and Tropical Medicine
Myles Allen, University of Oxford

Summary

- Currently, floods are an important problem in the UK. Floods are associated with few direct deaths, but the full effect on health, in terms of indirect mortality and morbidity due to infectious disease, mental health, and injuries is not known.
- Current evidence indicates that certain types of floods, such as Spring floods due to snow melt, are likely to become less frequent under climate change, while others, such as those directly linked to sustained Autumn and Winter precipitation, are likely to become more frequent. Risks will evolve differently in different regions and river catchments.
- There is good evidence to expect an increase in the frequency of heavy precipitation, with the greatest increases in frequency occurring for short duration, high-intensity events. Hence, an increased risk of flooding may accompany no change, or even a decrease in mean rainfall or in the duration of wet spells.
- Along the North Sea Coast there is a low risk of a very severe flood-event (as in 1953), but this risk will increase due to the rising sea levels unless coastal defences are upgraded appropriately.
- The direct health effects due to changes in windstorm intensity or frequency are likely to be small relative to other health effects of climate change in the UK.

Recommendations

- Implementation of flood health risk assessments
- Improved implementation of interventions to reduce health impacts associated with flood events, including the better use of current measures, such as guidelines for dealing with chemical contamination of flood waters, and post-flood support and counselling.
- Strengthening emergency planning and preparedness.

2.1 Introduction

Flooding and windstorms are a regular occurrence in the UK. Flash floods in Boscastle, Cornwall destroyed homes and businesses and swept away cars in August 2004. Heavy rainfall in early January 2005 also caused extensive, and in places repeated flooding across the North of England and Wales. In January 2005, a severe gale across Northern Ireland and North-West Scotland, with wind speeds in excess of 120mph, resulted in seven fatalities and loss of power to 85,000 households. The contribution of climate change to our current flood and windstorm risk has yet to be definitively quantified. Evidence has been strengthened, however, of the threat that climate change presents to future flood risk in the UK, and the consequent implications for human health.

The first part of this chapter will review advances in understanding the health impacts of flooding and windstorm events, while the second part will discuss the physical and model-based reasons why we may expect the risk of such events to be changing.

2.2 The health effects of flooding and windstorms

Since the last report for the Department of Health (2001), little new research on the health effects of flooding has been completed in the UK. Two reviews of the epidemiological literature have described the range of health effects: drownings, injuries, mental illness, and infectious diseases (Hajat *et al*, 2003; Ahern *et al*, 2005). A study following the flooding Lewes in 2002 found that people who were flooded experience a range of illnesses (Reacher *et al*, 2004). However, important knowledge gaps remain regarding the impact of flood events on long-term anxiety and depression, and indirect effects on mortality and use of health services in the 12 months following a flood event. Qualitative studies following flood events in Oxfordshire have shown that impacts on quality of life can be severe and long-lasting (Ohl and Tapsell, 2000; Tapsell *et al*, 2002).

Deaths

The 2001 report reviewed coastal and riverine flood risk separately. The health implications of these events are essentially the same. One of the most important factors for the risk of drowning is the speed of onset of the flooding, and the presence or not of some kind of warning.

Deaths associated with flooding in the UK are relatively few compared with other European countries. Only 3 deaths have been reported since 2001 (Table 2.1). The Environment Agency and Defra have no formal criteria for a flood death, but collate the information from situation reports that are completed for each flood. It is likely that official flood deaths are limited to persons who are swept away or drown in or near their own homes.

Recent reviews have confirmed that in industrialised countries, such as the UK, flood deaths are primarily associated with flash floods (Jonkman and Kelman, 2005), and are often vehicle-related (French *et al*, 1983). A review by Kelman (2003) also identified several other types of drownings associated with floods, e.g. persons having drowned after entering swollen rivers. Risk of death is also apparent from electrocution and carbon monoxide poisoning in the clean-up stages after the flood. [More than 20 episodes of electrocution and 5 deaths from carbon monoxide poisoning were reported after Hurricane Katrina in the US in 2005 (CDC, 2005).

Table 2.1. Major floods in UK since January 2000 and associated deaths

Area	When	No of households flooded	Reported deaths
Central & Southern England	New Year floods, 2002-2003	1029	1
Wales	February 2004	150	none
Boscastle, Cornwall	August 2004	60	none
Carlisle	January 2005	3000	3
River Tyne	Jan/Feb 2005	100	none

Source: Flood Risk Management, Environment Agency.

The Royal Society for the Prevention of Accidents (RoSPA) reports drownings per year associated with rivers or streams, and some drowning deaths attributed to flooding and high tides (RoSPA, 2005). Such numbers of drownings are likely to be incomplete as the “cause” will not be systematically reported. Defra and the Environment Agency do not collate national statistics on flood deaths or injuries, and, as discussed above, the definition of what constitutes a flood death is unclear and narrowly interpreted.

Age and gender related information on flood deaths is also incomplete. Published reviews have shown that men are much more at risk of drowning than women, probably due to more risky/heroic behaviour. Elderly people are particularly at risk of drowning in their own homes. For example, in the Carlisle floods two of the three deaths were in elderly women in the same street. In the 1953 ‘Big Flood’ in East Anglia, the majority of the deaths were in elderly persons (Baxter, 2005). Small scale flooding can also claim lives: an elderly person drowned in a basement flat in North London during an episode of very heavy rainfall in 2002.

Chemical hazards

The Health Protection Agency (HPA) Chemical Hazards and Poisons London Regional Unit has reviewed the implication of flooding for chemical hazards (Euripidou and Murray, 2004). There are risks to health from the contamination of flood waters with chemical waste, oil, diesel, pesticides, fertilisers, etc.. Flooding may lead to mobilization of dangerous chemicals from storage or remobilization of chemicals already in the environment, e.g. pesticides. A case study of heavy metal soil contamination after the flooding of the river Meuse during the winter of 1993-1994 concluded there was a potential health risk for river-bank inhabitants as a consequence of lead and cadmium contamination of the soil (Albering *et al*, 1999). Hazards may be greater when industrial or agricultural land adjoining residential land is affected. The UK has a high population density, with people living close to industrial installations (Young *et al*, 2004).

Mental health

Effects on mental health are now recognised as an important health consequence of flooding (Ohl and Tapsell, 2000). A systematic review of the epidemiological literature found that studies, mostly from the US, did provide good evidence of long term (> 6 months post flood) mental health complications (Ahern *et al*, 2005). A study in an elderly population affected by the 1993 US midwest floods, found that flooding increased the incidence of common mental disorders, even after adjusting for pre-flood morbidity

(Ginexi *et al*, 2000). A case-control study from the UK, found a four-fold increase in psychological distress among adults whose homes were flooded compared with those whose homes were not flooded (Reacher *et al*, 2004).

Infectious disease

Infectious disease risk concerns relate to contact with raw human or animal sewage/effluent. The Health Protection Agency (HPA) provides guidance advising against the consumption of food that has been exposed to flood water. Flood water itself is unfit for drinking and notices advising that tap water should be boiled are issued as necessary. Enhanced surveillance during floods for gastro-intestinal illness has revealed no apparent increases that can be attributed to recent flood events. However, an increase in self-reported illnesses is apparent following the floods in Lewes: skin disease, respiratory disease and stomach upsets (Reacher *et al*, 2004). The role of heavy rainfall in outbreaks of water borne disease is discussed in more detail in Chapter 5.

Effect of flooding on health service delivery

Flood events may also indirectly affect health by causing damage to hospitals, clinics and general practices. The siting of hospitals, care homes, schools, nurseries, and clinics in flood risk areas should be assessed. In Lewes, one general practitioner's office was flooded, and most of the patients notes were destroyed.

Floods in Dresden, Germany, in 2002 left large parts of the city without power and freshwater for several days (Meusel and Kirch, 2005). Four of the six main hospitals in Dresden were sited near the River Elbe and affected by the flooding. Two of the hospitals were evacuated.

Deaths and injuries from windstorms.

Deaths from windstorms are not collated by the Environment Agency. Information is available from a range of press reports. Many deaths are vehicle related, and a small but significant number are in outdoor workers.

Some recent deaths attributable to high winds:

- Four men died when they fell off a gantry while working on the Avonmouth Bridge in high winds (1999).
- In January 2005, a severe gale across Northern Ireland and North-west Scotland, with wind speeds in excess of 120mph, resulted in seven fatalities and the loss of power to 85,000 households.

A retrospective study of the injuries caused by high wind speeds during the storms of early 1990 found that that high wind speeds were associated with an increased risk of injury, especially when wind gusts exceeded 60 knots (Cugnoni and Whitworth, 2002). The apparent wind speed threshold, above which injury is more likely, corresponds to the curve used by the Met Office to predict wind-related structural damage.

2.3 Climate change, riverine flooding and extreme precipitation

The relationship between precipitation and flood risk depends critically on regional hydrology. There is no single type of extreme precipitation event that is relevant to flood risk. The floods of Autumn 2000 and January 2003 were associated with sustained (of the order 30-days or longer) catchment-scale rainfall anomalies that resulted in a gradual increase in water table levels as precipitation exceeded run-off capacity. In

contrast, the Boscastle flood in August 2004 was associated with a localised, slow-moving thunderstorm that resulted in over 180 mm of rainfall in under 5 hours over a single, relatively small river catchment area. Summer floods tend to be associated with short-duration extreme precipitation, but in other seasons all time-scales are relevant (for example, the Carlisle floods in 2005 were associated with a specific storm system).


Our understanding of the influence of climate change on large-scale, sustained precipitation anomalies is further advanced than our understanding of short-term, localised anomalies. On physical grounds, it could be argued that we should expect more severe thunderstorms in a warmer, moister atmosphere, but since many factors affect the intensity and distribution of thunderstorms, a simple link between temperatures and thunderstorm-related flood risk cannot be expected.

Modelling studies of short-term extreme precipitation show a tendency for greater fractional increases in intensity at higher return times: that is, the percentage increase in precipitation events that are expected to occur once every 30 years is greater than the percentage increase in events that are expected to occur once per year (see chapter 1). This is illustrated in Figure 2.1 (adapted from Allen and Ingram, 2002), using data from the HadCM3 climate model. The green dashed line in the upper panel shows the intensity, in millimetres per day, of rainfall events as a function of their return time (the reciprocal of the number expected in a given year) in the model control climate, while the blue solid line shows how these intensities change by years 2070-2100 in a simulation of the response to the A2 emissions scenario (the medium-high scenario). The red dotted line shows the fractional increase (how much the blue curve is higher than the green one): for this particular model, scenario and grid-box, there is a tendency for the fractional increase in the intensity of frequent events (e.g. 0.1-year return time, or 10 events per year) to be less than the fractional increase in rarer events (e.g. 3-year return time).

If we examine all grid-boxes at the same latitude (lower panel), a noisier picture emerges. Nevertheless, there is still a significant shift upwards in the distribution of increases as we move from frequent (10 per year) to infrequent (one in 10 year) events. The apparent downturn towards the most infrequent events is almost certainly a sampling artefact, since the simulations in question will have contained very few events of these magnitudes.

This overall result is confirmed by assessments using higher resolution regional climate models, as described in Chapter 1 on Climate. Changes in precipitation tend to be concentrated in high-intensity, short-duration events, so no change or even a decrease in seasonal mean rainfall may be accompanied by an increase in flash flood risk. Hence the *decline* in the length of extreme wet spells reported in Chapter 1 is, somewhat counter-intuitively, consistent with an *increase* in the risk of floods associated with short-duration precipitation events.


Figure 2.1 Intensity of daily precipitation events as a function of return time for the HadCM3 climate model, comparing control climate (top panel, green dashed line) with years 2070-2100 of a run driven with the SRES A2 forcing scenario (top panel, blue solid line). Red dotted lines and right-hand axes show percentage increase in intensity as a function of return time (figure courtesy of Pardeep Pall, University of Oxford, based on data provided by Jonathan Gregory, Met Office).



When we consider longer duration precipitation events, such as those responsible for the Autumn 2000 and Winter 2002/2003 floods, the picture becomes more complicated and results more model-dependent. Figure 2.2 (left panel) shows, purely as an illustration, how the intensity of 30-day extreme precipitation events vary in the grid-boxes corresponding to the Lewes catchment area (which suffered severe flooding in Autumn 2000) as greenhouse gases are increased from 1860 conditions (solid line), through 2000 (heavy dotted line) to 2090 (dashed line), following a simple 1%/year increase from 1990⁴. In this particular model (HadRM2 nested into the HadCM2 global climate model: see Huntingford *et al*, 2003, for further details) we see that a monthly average rainfall of 6.7 mm/day would occur on average once in 10 years in 1860, while a monthly average of 8.3 mm/day would occur with this frequency in 2090. Equivalently, a 30-year event in 1860 becomes a 12-year event in 2000, and a 4-year event in 2090. As in Figure 1, the fractional increase in intensity increases for rarer events, with a 13% increase between 1860-2090 for 1-year events, increasing to a 23% increase for 10-year events.

⁴ Note that this represents a stronger forcing than that used in figure 1

Figure 2.2. Left panel: Magnitude (in millimetres per day for 30-day average rainfall) of precipitation events as a function of return time in 1860 (solid line), 2000 (heavy dotted line) and 2090 (dashed line) in the HadRM2 regional climate model points corresponding to the Lewes catchment, forced with observed greenhouse gas increase to 1990 followed by 1%/year to 2100.



The light dotted lines in the left panel of Figure 2.2 show how a simple representation of uncertainty in the global mean response, following Allen *et al* (2000), can provide a range of possible changes in the distribution of extreme precipitation that are attributable to past human influence on climate. The light dotted lines are all lower than the heavy dotted line because all estimates of the net anthropogenic climate change to date are lower than the estimated greenhouse-only warming simulated by the HadCM2 climate model, primarily because the net effect of non-greenhouse anthropogenic drivers of climate change is almost certainly leads to a reduction in ambient temperatures.

Allowing for uncertainty in the response provides a distribution of possible changes in risk that are consistent with data available to date: this is shown in the right hand panel. The top axis shows the fraction by which an occurrence of monthly mean precipitation exceeding 7.5mm/day (a 40-year event in the 1860 climate of this model, light horizontal line in the left hand panel) is estimated to have increased due to past human influence on climate. The best estimate (peak of the distribution) corresponds to a doubling of risk, with some uncertainty on either side. Should such an event occur, the bottom axis shows what fraction of the risk is attributable to, or to what extent it can be “blamed on”, the influence of previous human activities on climate. (See Stott *et al*, 2004, for a description of the application of the epidemiological concept of attributable risk to weather events and an application to the 2003 European summer heatwave).

We emphasise that these results are presented simply to illustrate the methodological progress that has been made since the previous report. Results remain too model-dependent for quantitative conclusions to be drawn regarding risks in the real world: a follow-up study using the HadRM3H regional climate model found results consistent with no increase or even a reduction in the risk of extreme 30-day precipitation events. Hence, uncertainty exists in modelling at both a global and a regional side and both should be considered.

2.4 Climate change and windstorms

Evolving windstorm risk in the UK has been comprehensively assessed using data from the UKCIP02 scenarios (Hansen *et al*, 2004). This study focussed on windstorm risk due

to deep cyclonic depressions⁵ that are responsible for the bulk of windstorm damage in the UK, primarily occurring in the autumn, winter and spring seasons. High winds associated with thunderstorms, squall lines or tornados would not be resolved by current models.

Reflecting the modest changes in cyclone activity in the regional climate model, changes in maximum daily wind speed and numbers of days with gales are also small and spatially variable, with small reductions projected in some regions and small increases in others. Any projected change in large-scale (cyclonic) windstorm activity is much smaller than the minimum estimated uncertainty in the projections, leading us to conclude, on the basis of evidence available to date, that climate change will not significantly effect health via changes in windstorms.

An important caveat to this conclusion is that windstorm risk is known to be closely associated with the value of the North Atlantic Oscillation (NAO), and while the evidence is ambiguous, there are some indications that current climate models are underestimating the response of the NAO to anthropogenic climate change (Gillett *et al*, 2003). If the next generation of models project larger changes in the NAO, we can expect correspondingly larger changes in windstorm risk.

2.5 Sea level rise, coastal flooding and storm surges

Coastal flooding and storm surge risks respond both to changes in windstorm activity and to the background sea level. The 1953 Big Flood, which killed 307 people, was caused by a combination of a windstorm in the Atlantic that caused a major storm surge at the same time as a very high tide.

In a study using the HadRM2 regional climate model, which generally simulated much larger changes in cyclonic activity than HadRM3, similar contributions from both factors were found to increase storm surge risks around the British Isles (Flather and Smith, 1998). With the smaller changes in windstorm activity predicted by HadRM3, the relative contribution of background sea level rise is higher. While the considerable uncertainties in windstorm activity were noted in the previous section, sea level rise (particularly that due to thermal expansion of the oceans) is much more predictable. Hence we can predict with some confidence that the risk of coastal flooding will increase due to anthropogenic climate change over the 21st Century.

Projected changes in storm surge risk are highly variable between locations, but can be substantial. The frequency of a 1-in-500 year extreme water level at Immingham on the Northeast coast was projected to increase to once every 15 years. The relative contributions of windstorm versus background sea level rise also vary between regions. Windstorm activity may dominate storm surge risk in Scotland, while both windstorm and background sea level rise contribute to the flood risk in East Anglia (Tsimplis *et al*, 2005).

The distinction between changes in storm surge risk due to windstorms versus background sea level is important because of their different relative predictability. The actual flood risk of an increase in background sea level is likely to be small provided this (highly predictable) factor is taken into account in the design of coastal defences, sewage systems and other susceptible infrastructure.

⁵ A depression was defined as a coherent synoptic system with a minimum central pressure $\leq 1,000$ HPa, while a deep depression had a minimum central pressure ≤ 970 HPa and/or a deepening rate ≥ 9.5 HPa in 12 hours.

2.6 Populations at risk of flooding

Many factors affect the flood risk for a given catchment, and climate changes represent only one part of this risk. The Environment Agency estimates that currently around 5 million people in 2 million properties live in flood risk areas in England and Wales. DEFRA estimates that 80,000 properties are at risk in towns and cities from flooding caused by heavy downpours that overwhelm urban drains – so-called ‘intra-urban’ flooding.

Sea level rise and tidal surge flooding are already threatening coastal lowlands, for example around The Wash, Norfolk, Suffolk and Humber. The Thames Barrier protects the 150km² of London that lies below the high tide level. It was closed 55 times to protect London from tidal surges between 1983 and 2004, and more than 50% of these occasions have occurred during the last five years. Of flood plains in the UK that of the Humber river has the greatest concentration of people – apart from that of the Thames. The tidal barrier in Hull and other flood defences protect over 300,000 people in the area.

Flood risk is not distributed equally among the population in terms of socio-economic status. In England, the people at greatest risk of tidal flooding are those on lower incomes, particularly in East Anglia. However, a spatial analysis that linked the population identified by the Environment Agency’s indicative-floodplain-maps with census data has shown inconsistent results (Fielding and Burningham, 2005).

An important consideration is *Residual Flood Risk*. Where there is a high standard of defence, the probability of flooding is low, but the consequences in terms of damage and threats to life, should flooding occur, are enormous. An estimated 1.25 million people live and work in the defended Thames tidal floodplain.

Box 2.1 Foresight Flood Project – main findings (Foresight Report, 2001)

- Climate change is an important factor in increasing flood risk, particularly through the impacts of rising sea levels and more stormy weather.
- Other important factors include the way we use land, increased urban development and the effects of increased wealth and higher standards of living.
- The number of people at a high risk from flooding could rise from 1.5 million to 3.5 million by 2100.
- More effective land management will help reduce the risks in most scenarios. However, in the worst case scenario these are of little benefit and greater use of flood defences and coastal re-alignment will be required.

The Foresight Flood Project did not estimate future flood deaths or costs attributable to health effects.

Acknowledgements

The authors are grateful to Marilyn Mitchell at the Environment Agency for her involvement with this work.

References

- Ahern, M. J., Kovats, R. S. Wilkinson, P. Few, R. and Matthies, F. (2005) Global Health impacts of floods: epidemiological evidence. *Epidemiological Reviews*, **27**, 36-46.
- Albering, H. J., van Leusen, S. M. Hoogewerff, J. A. and Kleinjans, J. C. S. (1999) Human health risk assessment: a case study involving heavy metal soil contamination after the flooding of the river Meuse during the winter of 1993-1994. *Environmental Health Perspectives*, **107**, 37-43.
- Allen, M. R., Stott, P. A., Mitchell, J. F. B., Schnur, R. and Delworth, T. (2000) Quantifying the uncertainty in forecasts of anthropogenic climate change. *Nature*, **407**, 617-620.
- Allen, M. R. and Ingram, W. (2002) Constraints on future changes in climate and the hydrologic cycle. *Nature* **419**, 224-232.
- Baxter, P. J. (2005) The east coast Big Flood, 31 January - 1 February 1953: a summary of a human disaster. *Philosophical Transactions of the Royal Society: London series A*, **363**, 1293-1312.
- CDC (2005) Carbon monoxide poisoning after hurricane Katrina--Alabama, Louisiana, and Mississippi, August-September 2005. *Mortality and Morbidity Weekly Report*, **54**, 39, 996-998.
- Cugnoni, H. L. and Whitworth, I. (2002) Injuries related to wind speed. *Ann.R.Coll.Surg.Engl.* **74**, 4: 294-296.
- Euripidou, E. and Murray, V. (2004) Public health impacts of floods and chemical contamination. *Journal of Public Health* **26**; 4: 376-383.
- Fielding, J., and K. Burningham, 2005, Environmental inequality and flood hazard: Local Environment, v. 10, no. 4, p. 379-395.
- Flather, R. A., and Smith, J. A. (1998) First estimates of changes in extreme storm surge elevations due to the doubling of CO₂. *The Global Atmosphere and Ocean System* **6**, 193-208.
- French, J., Ing, R., von Allmen, S. and Wood, R. (1983) Mortality from flash floods: a review of national weather service reports, 1969-81. *Public Health Rep.* **98**, 584-588.
- Gillett, N. P., Graf, H. F. and Osborn, T. (2003) Climate change and the North Atlantic Oscillation, in JW Hurrell, Y Kushnir, G Ottersen, and M Visbeck (eds.), North Atlantic Oscillation: climatic significance and environmental impact: American Geophysical Union. pp. 193-209.
- Ginexi, E. M., Weihs, K., Simmens, S. J. and Hoyt, D. R. (2000) Natural disaster and depression: a prospective investigation of reactions to the 1993 midwest floods. *Am.J.Community Psychol.* **28**, 4, 495-518.
- Hajat, S., Ebi, K. L., Kovats, R. S., Menne, B., Edwards, S. and Haines, A. (2003) The human health consequences of flooding in Europe and the implications for public health: a review of the evidence. *Applied Environmental Science and Public Health*, **1**, 1, 13-21.
- Hansen, C., Holt, T. and Palutikof, J. P. (2004) An Integrated Assessment of the Potential for Change in Storm Activity over Europe: Implications for Insurance and Forestry in the UK. Technical Report **12**. Norwich, Tyndall Centre.
- Huntingford, C., Jones, R. G., Prudhomme, C., Lamb, R., Gash, J. H. C., and Jones, D. A. (2003) Regional climate-model predictions of extreme rainfall for a changing climate. *Q.J.R. Meteorol.Soc.* **129**, 1607-1621.
- Jonkman, S. N. and Kelman, I. (2005) An analysis of causes and circumstances of flood disaster deaths. *Disasters* **29**, **1**, 75-97.
- Meusel, D. and Kirch, W. (2005) Lessons to be learned from the 2002 floods in Dresden, Germany, in W Kirch, B Menne, and R Bertolini (eds.) *Extreme weather events and Public Health Responses: Springer-Verlag*, p. 175-184.
- Ohl, C. A. and Tapsell, S. (2000) Flooding and Human Health. *British Medical Journal* **321**, 11 Nov, 1167-1168.
- Reacher, M. et al, (2004) Health impacts of flooding in Lewes: a comparison of reported gastrointestinal and other illness and mental health in flooded and non flooded households. *Communicable Disease and Public Health* **7**, 1, 1-8.

- RoSPA (2005) Drowning statistics in the UK: 2002, Royal Society for the Prevention of Accidents,
<http://www.rosipa.com/waterandleisuresafety/drownings/2002statistics.htm>
- Stott, P. A., Stone, D. A. and Allen, M. R. (2004) Human contribution to the European heatwave of 2003. *Nature*, **432**, 610-614.
- Tapsell, S., Penning-Rowsell, E., Tunstall, S. M. and Wilson, T. M. (2002) Vulnerability to flooding: health and social dimensions. *Philosophical Transactions of the Royal Society: London series A*, **360**, 1511-1525.
- Tsimplis, M., Woolf, D., Osborn, T., Wakelin, S., Wolf, J., Flather, R. A., Shaw, A. G. P., Woodworth, P. and Challenor, P (2005) Towards a vulnerability assessment of the UK and Northern European coasts: the role of regional climate variability. *Philosophical Transactions of the Royal Society* **363**, 1329-1358.

3 Vector borne diseases and climate change

(See also section 4.3 of 2001 report)

David Rogers, University of Oxford
Sarah Randolph, University of Oxford
Steve Lindsay, Durham University
Stephen G. Willis, Durham University

Summary

- Health authorities need to remain alert to the possibility of future European malaria outbreaks or to the arrival in the UK of better European vectors of malaria.
- Any malaria outbreaks in the UK, however, are likely to be rare and on a small scale involving a small number of people. Prompt reaction to any outbreaks will reduce the chances of endemic malaria transmission in the UK.
- Any increase in tick-infested areas is more likely to be the effect of changing agricultural and wildlife management practices than the effect of changes in climate alone
- People engaging in leisure activities are most likely to contact Lyme disease infected ticks during spring and autumn, when ticks are likely to be most active. Whether, under various scenarios for climate change, there will be an increase in numbers of infected ticks at these times is still uncertain.
- None of the predicted climatic changes suggest that the UK will be threatened by the Tick-borne Encephalitis. Overall there may be a marked decrease in the extent of this pathogen, although some areas free at present may be invaded in continental Europe.
- A range of other insect-related problems may change with the predicted climate changes in the UK.

Recommendations

- Monitor changing risks as they are happening, both within the UK and to holiday makers.
- Examine the impact of multivariable environments on the spatial and temporal patterns of disease distribution and intensity for both temperate and tropical vector-borne disease systems.
- Establish a surveillance system for monitoring the distribution and abundance of arthropod vectors in the UK.

3.1 Introduction

Diseases caused by vector-borne pathogens are the result of complex interactions between three very different sorts of organisms:

- the vertebrate host,
- the invertebrate (usually arthropod) vector,
- the parasite (viruses, bacteria, etc.).

Each partner in the triangle is responsive to environmental changes in ways that make it very difficult to predict the altered outcome of these interactions. Rates of the biological processes (e.g. birth, death, development, biting by blood-sucking vectors, transmission between host and vector) vary independently and often in opposite directions in response to each climatic variable. Increased temperature, for example, generally accelerates development of both vectors and pathogens within those vectors. This causes vectors to bite more frequently (increasing transmission rates), but may also cause higher vector mortality rates (which decreases transmission). Increased dryness may exacerbate vector mortality, and additionally, decreases the availability of breeding sites for vectors such as mosquitoes. Although temperature and dryness may vary together, they will do so to different extents in different regions. In temperate regions with markedly seasonal climates, indices of mean annual climate changes alone are not sufficient for predictions of the risk of infection in the human population. For example, no matter how suitable most of the year may become in the United Kingdom for sub-tropical vectors, either a hot dry summer or a cold winter may act as a bottle-neck, reducing vector survival.

We therefore urge caution in drawing conclusions about the future incidence of infection from existing relationships between single climatic variables and disease. In general, more than a single variable will determine transmission rates, and correlations observed today may fall apart, or at least change shape, if the underlying biological processes are disrupted by climate change beyond the normal fluctuations upon which those correlations are based.

3.2 Methodology

Baseline and modelled climatologies

The CRU 50km British Isles climatology and 0.5° Global climatology (UKCIP02 Scenarios based on the HadRM3H regional model, Chapter 1) were used to interpret and model present-day distributions of vector-borne diseases. To these surfaces were added the HadRM3H model (difference) predictions for the ‘medium-low’ and ‘medium-high’ scenarios. These differences were also scaled up or down by factors derived from UKCIP98 Appendix 6 (UKCIP98 section, ΔT columns) to produce the equivalent ‘high’ and ‘low’ global surfaces for the 2020, 2050 and 2080 periods. The same correction factors were applied to all climate variables (i.e. including vapour pressure and precipitation), following current accepted practice (M.Hulme *pers. comm.*). Thus the future scenarios are a combination of the observed 1961-1990 climatology and the predicted GCM differences under the variety of scenarios considered: this, rather than using both modelled 1961-90 climatology and difference information, also follows standard practice (M. Hulme *pers. comm.*).

Before using any of the future scenario information the HadCM2 monthly imagery had to be re-sampled to the same image dimensions as the 0.5° global datasets. This was done using cubic spline interpolation (by columns and then by rows).

Modelling disease distributions

There are a number of options for modelling the impact of future climates on vector-borne and other transmissible diseases. This section discusses some of the alternatives:

a) The Biological approach

All biological processes are temperature sensitive, but the impact of this sensitivity on disease transmission is often difficult to predict (see section 3.1). The temperature-dependence of a number of important variables has been established experimentally, and may be included in predictive transmission models for a warmer world. Arising from this, developmental processes are often modelled by day-degree summation, where temperature differences from a developmental threshold are summed over time until a total is reached that represents the total developmental requirements, at which point development is judged to be complete. Such temperature summation may be used to model the incubation period of a parasite within vector insects, thus providing another input into general models for disease transmission.

- The **biological approach** is used to predict the impact of the various GCM scenarios on the potential for transmission of *vivax malaria* in different parts of the UK.

Whilst biological modelling of this sort is the ideal solution to the problem of predicting the impact of future climates on disease transmission, at present there are insufficient data to use such models to predict how mosquito abundance (an important component of these models) will change under different scenarios of climate change. Warming conditions from a cold temperature start are likely to increase vector numbers, but from a warm temperature start may well decrease numbers. Such population changes are ignored in all existing models of this sort, and the results of model predictions of future risks relative to those of the present, assume constant vector populations. A further limitation is that the models tend to be driven only by temperature variables; the influence of all other variables is ignored. Increasing or decreasing moisture, in concert with temperature changes, is a critical factor that should also be considered, but is difficult to assess.

Methods such as this, and the following ones, make predictions about the changing potential risk from biting vectors. Whether or not this potential risk is realised depends on the human response to varying biting frequencies; for example, while a low biting rate may be ignored, higher rates may elicit avoidance behaviour which reduces human-vector contact.

b) The Statistical approach

In this second approach, the present-day distributions of vectors or diseases of potential importance are statistically matched to current climatic variables, to provide a multi-variate description of present-day areas of disease risk. This understanding is then applied to future scenarios, and future distributions are predicted for them. The result of climate change might be that tropical vectors and the diseases they transmit may arrive in Northern Europe and the statistical approach gives some indication of the probability of this occurring. The approach, therefore, is essentially a 'pattern-matching' exercise, from which conclusions may be drawn about the likely climatic sensitivity of vectors and/or diseases. Obtaining a good fit of the present-day distributions to present day climates is a necessary first step in this modelling exercise. Distributions are modelled here using maximum likelihood methods and discriminant analytical models with step-

wise inclusion of temperature, vapour pressure and precipitation variables. Before analysis, all climate surfaces were temporally Fourier smoothed (Rogers *et al*, 1996) and the means, maxima and minima of the smoothed variables were used in the analysis. The step-wise inclusion method identifies sequentially those variables most important in distinguishing presence and absence areas: we imagine that this reflects their biological importance (although this assumption urgently needs to be tested). For the present report, the top five selected variables only were used to make the predictive maps. Climate matching is well able to capture the rather subtle interactions between predictor variables (many of which co-vary) that are often important in shaping distributions.

- The **statistical or pattern-matching approach** is illustrated for the cases of **global malaria** and **tick-borne encephalitis (TBE) in Europe**. The changing distributions of both diseases are modelled for a variety of future scenarios. Changing risks, both to the UK population directly and to UK travellers to various regions, are predicted.

The drawback of the pattern-matching approach is that it is essentially a statistical inference method, based on the past, that may not be a reliable guide to distributions in a climatically changed world where co-variation between climate variables may be different. It is also possible that the variable selection method will identify biologically spurious variables: it is therefore important to select the variables submitted to the analysis on the basis of prior biological understanding.

c) The Climate Envelope approach

This is a version of the above approach that matches current UK climates to other parts of the world and then examines what diseases are prevalent in them. Presumably there is some risk that the same diseases could affect the UK depending also upon relative standards of living in the matched countries. The risks of future diseases may then be investigated by matching *future* UK climates to *present* global climates, to answer the question 'Where, at the present time, are climates similar to those that the UK will experience in 20, 50 or 80 years' time?'. The diseases prevalent in such areas may be those we need to guard against.

- The **climate envelope approach** is applied to **current** and **future UK climates**. It is the only approach possible for situations where we lack any biological information for process-based modelling, or disease or vector distribution maps for the statistical approach.

The interpretation of the results from the climate envelope approach involves substantial guesswork, not only because disease situations in other countries are even less adequately known than in the UK (with the notable exception of the USA), but also because the ways in which people live within their environments vary in so many ways. For example, if future UK climates match those of Southern France today, what are the future risks in the UK of leishmaniasis, a protozoal disease mostly of dogs in the Mediterranean region, but occasionally transmitted by the sand-fly vectors to humans?

d) The Expert Opinion approach

When the above approaches cannot be applied, we must rely on expert opinion, often notable by its absence, its contradictions, or its inaccuracy. This in reality reflects the poor quantitative understanding we have of virtually all vector-borne and similar diseases - an ignorance that affects all prediction attempts in one way or another.

Currently the geographically variable importance of Lyme disease (O'Connell *et al*, 1998) reflects either biological reality, or inadequate diagnosis: increasing recognition in the UK has resulted in increased numbers of reported cases here (Cannell *et al*, 1993) which probably has nothing to do with climate change. When future predictions are made of

diseases such as this, already present in the UK, we fall back on a combination of biological insight to suggest how disease prevalence may change in vector populations, and guessed predictions of how altered British leisure activities in the future warmer world will affect human interactions with these infected vectors.

- The **expert opinion approach** is used to make predictions about the future importance of **Lyme disease**, caused by bacteria transmitted by ticks. Lyme disease already occurs in the United Kingdom, throughout the rest of Europe and into the far East of the Russian Federation.

A number of other potential future problems for the UK can be identified using expert opinion on the likely increase of insects and other creatures that are known to generate nuisance or public health problems at the present time. Such problems often come to the attention of experts only from articles in the local or national press, or from talking with colleagues whose advice has been sought by the affected regional health authorities. Neither is a particularly reliable or consistent measure of the real extent of such problems. Finally, expert opinion may be used to second-guess the future arrival in the UK of migrant pests such as locusts, or of imported pests such as the Asian tiger mosquito that has already spread rapidly in the USA and large parts of Europe.

- **Expert opinion** is used in this report to produce a list of potential problems in the UK from outbreaks of **urticaceous caterpillars, stinging insects or nuisance flies** in houses or around kitchens, to more direct threats such as local outbreaks of **blood-sucking black-flies** (*Simulium* spp.).

3.3 Malaria

Malaria is one of the world's most important infectious diseases killing 742,000 children under five years old every year (Snow *et al*, 2003). The disease is transmitted by inoculation of the parasite during feeding by certain Anopheline mosquitoes, which breed in both fresh and brackish water. Despite attempts to control this disease there are between 300-660 million clinical cases each year and this number may be rising (Snow *et al*, 2005). Of the four species of human malaria, *Plasmodium falciparum* is the most lethal and is widespread throughout the tropics. *P. vivax* is less harmful, but is still responsible for much illness and occurs widely in the tropics, although it is uncommon in much of Africa. The problem of malaria is particularly worrying because of the rapid spread of drug-resistant strains of the parasite and the possibility of untreatable forms of malaria.

Domestic malaria (predicted using the biological approach)

Malaria was once common in many marsh communities in Southern England between the 16th and 19th centuries (Dobson, 1997) and some indigenous malaria occurred at the beginning of the 20th Century. Those areas most badly affected included the Fens, Thames estuary, South-East Kent, the Somerset levels, the Severn Estuary and the Holderness of Yorkshire (Shute and Maryon, 1974). Malaria declined progressively from the 1820s onwards due to a number of factors. Drainage schemes in the marshlands shrank mosquito-breeding sites. Housing improved and became less suitable for resting mosquitoes, which prefer damp and dark quarters. People began to sleep in separate rooms, often upstairs, making it more difficult for a mosquito to locate a human blood meal. Cattle numbers rose and were stabled away from homes, providing an alternative source of blood and reducing the chances of malaria transmission. At the same time improvements in medical practice occurred and quinine, an effective anti-malarial, became more affordable (Newman, 1919).

In 1917 and 1918 there were around 330 cases of locally-transmitted *vivax* malaria when infected servicemen returning from overseas were billeted near salt marshes on the Thames Estuary (James, 1920). After that, effective control was achieved by making malaria a notifiable disease, with appropriate swift treatment and control action. All reported cases of indigenous malaria this century were *vivax* malaria, except for one unusual case of *falciparum* malaria in Liverpool.

Mosquito vectors and malaria transmission

There are six species of Anophelines in Britain capable of transmitting both temperate and tropical strains of *vivax* malaria: *Anopheles algeriensis*, *An. atroparvus*, *An. claviger*, *An. daciae*, *An. messeae* and *An. plumbeus* (Linton *et al*, 2005). In the past British malaria was associated with salt marshes where *An. atroparvus* was common. The aquatic stages of this mosquito are found in fresh or brackish water and the adults occupy houses and feed readily on people. *An. atroparvus* can also transmit European strains of *P. falciparum*, but is completely refractory to strains of the same parasite from the tropics (Ramsdale and Coluzzi, 1975). This mosquito is therefore considered the most important potential vector of malaria in the UK. However, field studies on the Isle of Sheppey, the last place in England to experience a malaria epidemic, show that this mosquito is now relatively rare and is unlikely to bite large numbers of people. Another possible vector is *An. plumbeus* that has a widespread distribution, that includes London and its suburbs. It typically breeds in tree holes, although larvae have also been found in water held in old tyres (Karch, 1996). Recently it has been shown that *An. plumbeus* is capable of transmitting *P. falciparum* (Curtis 2003).

How likely is it that *vivax* or *falciparum* malaria could be transmitted in the UK? Malaria transmission requires that a potential vector should feed on someone carrying gametocytes, the stage of the parasite that is capable of maturing and becoming infective within a mosquito. Between 9-24 days after taking an infectious blood meal of gametocytes (depending on the temperature (Boyd, 1949), a vector mosquito will be able to transmit the infection to anyone that it bites.

In 2003 there were 206 cases of imported *vivax* malaria and approximately 66% of these were contracted by visitors to the Indian Subcontinent (HPA, 2004). Since there are relatively few *vivax* cases in the UK, and most of these are amongst people of Asian decent who tend to live in major urban areas, the possibility of *An. atroparvus*, a coastal mosquito, biting an infectious patient are remote. It is therefore extremely unlikely that *vivax* malaria will be locally transmitted within the UK.

With *falciparum* malaria, the risk assessment is not so straightforward. In 2003 there were 1,339 imported cases of *falciparum* malaria, of which 71% were from west Africa; mainly Nigerians and Ghanaians. Many of these people live in London (Williams *et al* 2002) and other urban areas in the South of England, areas where *An. plumbeus* occurs. An important question is are these mosquitoes likely to feed on patients carrying gametocytes? A person bitten by an infective mosquito will become ill after about 14 days (Gillies and Warrell, 1993) and then go on to produce gametocytes about 10 days later (Boyd, 1949). Since nearly 90% of *falciparum* cases are not detected until 1-5 months after arrival in the UK (HPA, 2004) it suggests that many will be carrying gametocytes in their bloodstream for several weeks. It is also the case that gametocytes may not be completely eliminated by chemotherapy (Targett *et al*, 2001). A further consideration is that the peak in imported malaria cases coincides with the period when *An. plumbeus* is most abundant (Williams *et al* 2002). We are not suggesting that *An. plumbeus* is responsible for local transmission since it is a relatively rare mosquito and will bite few people. We would therefore not expect more than a few very cases of autochthonous malaria in the UK over the next 50 years. Indeed one is more likely to be struck by lightning than to get malaria from an English mosquito.

Malaria and climate


P. vivax is better suited to the British climate than is *P. falciparum*. It requires lower temperatures (by 1-2°C) than *falciparum* to develop equally rapidly in mosquitoes, and thus does better at cooler temperatures. *Vivax* parasites, unlike *falciparum* parasites, also sequester in the liver of an infected person, and are later released to infect new generations of mosquitoes in the spring. As few parasites develop in mosquitoes below 15°C, the season for potential transmission in the UK is between June and September.

Temperature and rainfall both influence the level of malaria transmission (Lindsay and Birley, 1966). Higher temperatures increase the rates of mosquito development, female mosquito feeding and maturation of the malaria parasites within the mosquito, but may decrease adult mosquito survival. Rainwater provides mosquito breeding sites and a humid environment, conducive for vector survival.

Impact of temperature changes


Here the risk of *vivax* malaria in the UK is modelled for *An. atroparvus*. Maps of malaria suitability under recent climate (average 1961-90) and a range of future climate scenarios (Figures 3.1 and 3.2) show the number of months that *vivax* malaria, if it were introduced, could persist each year in different parts of the country. These maps are a rough guide of risk, based on the effects of changing temperature on variables in the transmission process, but they do not take into account changes in precipitation, humidity or the availability of mosquito breeding sites. These latter factors may not be so critical for the UK because the risk of malaria transmission is likely to be highest near extensive areas of wetland, which provide numerous breeding sites and are less affected by small changes in rainfall.

Figure 3.1 Present areas where the climate could support *vivax* malaria. Shading represents the number of months where *vivax* malaria could be transmitted (grey =0, blue =1, yellow =2, orange =3). Red circles show malaria cases in the 19th Century. Based on the UKCIP for 1961-90.



The present day distribution of malaria risk corresponds extremely well with past records of the distribution of malaria in England (Fig. 3.1) (Nuttall *et al*, 1901). Thus, we are confident that our temperature-malaria model is relatively robust. Under all climate-change scenarios, the risk of transmission is predicted to increase in the South of England, spreading Northwards to the Scottish Borders.

Figure 3.2. Projected future risk maps for malarial transmission under a medium-low (a, b and c) and a medium-high (d, e and f) climate change scenario. Maps show risk for the period 2020s (a and d), 2050s (b and e) and 2080s (c and f). Based on UKCIP02.



- At present, in only a few months in the South of the United Kingdom are temperature conditions permissive for transmission of *P. vivax* malarial parasites by indigenous vector mosquitoes. Although such transmission occurred in the historical past, it is a minor threat at present because living conditions have improved considerably since Shakespeare wrote of the 'ague' (malaria fever). If the climate becomes warmer, conditions for transmission become more favourable, and last for longer. It is likely that our present standards of living will ameliorate this increasing threat to a large extent, but not necessarily wholly in high risk areas.

Impact of habitat changes

The distribution of *An. atroparvus* is largely restricted to salt-marshes, because it breeds mainly in brackish water. At present there are 42,251 ha of salt-marsh in Britain, with the largest areas, 8,525 ha, along the Greater Thames Estuary in Essex and Kent (Davidson *et al*, 1991). Coastal wetlands are being reduced by drainage and other land 'improvements'. Rises in sea-level that breach sea defences and inundate lowlands that are at present prevented from adapting naturally to saltwater, may result in less salt-marsh. Elsewhere, gradual saltwater intrusion into coastal lowlands may increase breeding sites for *An. atroparvus*. With summer droughts, there is likely to be a decline in the numbers of *An. plumbeus* as tree holes dry out. In contrast, other mosquito species may find more breeding sites in pools left in river beds, and in water butts. There will be greater exposure to mosquitoes as people stay outdoors in warmer summer evenings, or sleep with the windows open.

It is possible that climate changes will allow new vector species to become established in Britain. This would be most serious if it involved better European vectors of *vivax*, such as *An. saccharovi*, *An. labbranchiae*, *An. superpictus* and *An. sergentii*.

- Health authorities need to remain alert to the possibility of future European malaria outbreaks, as in Italy after 40 years of being free of malaria (Simini, 1997), or to the arrival in the UK of better European vectors of malaria. Any malaria outbreaks in the UK, however, are likely to be rare and on a small scale involving a small number of people. Prompt reaction to any outbreaks will reduce the chances of endemic malaria transmission in the UK.


Travellers' malaria (predicted using the statistical approach)

There are two different threats posed by changes in the distribution of malaria in parts of the world other than the UK. First, aircraft travelling from malaria-endemic countries may bring infected mosquitoes into non-endemic countries. Such outbreaks occur occasionally around international airports. Between 1969 and 1996 there have been 60 such cases reported in Belgium, France, Germany, Italy, Netherlands, Spain, Switzerland and the UK (Danis *et al*, 1999). Between 1969 and 1999 there have been 14 cases of airport malaria in the UK (Gratz *et al*, 2000). In the summer of 2001, 52 aircraft arriving at Gatwick Airport from Africa were searched by experts for a total time of 43 hours (Hutchinson *et al*, 2005). This represented a search of about 25% of all flights that left Africa at night when one would expect the greatest number of mosquitoes to enter the aircraft. No malaria mosquitoes were found, demonstrating that the current practice of disinfection of aircraft travelling from the tropics to the UK is effective.

The second threat is of imported malaria, with its associated health system costs. There is no vaccination against malaria, and chemoprophylaxis is not guaranteed to be fully effective against a growing problem of multi-drug resistant parasites. Any increase in malaria endemic areas, alongside expanding jet travel, will raise the chances of travellers returning to the UK with malaria, including the far more dangerous forms found in the tropics. At present about 2,000 cases of travellers' malaria are reported each year to the

HPA Malaria Reference Laboratory (Figure 3.3). The recent fall in malaria cases probably reflects the reduction in travel that has occurred since 9/11. Of these, around 78% are infections with life-threatening *P. falciparum* (HPA, 2004). The numbers of *falciparum* cases have continued to rise over the last decade (Fig. 3), with 71 deaths from *falciparum* malaria occurring between 1998 and 2003.

Figure 3.3 Annual cases of malaria reported to the HPA Malaria Reference Laboratory, UK. (adapted from CDR Weekly, Vol 14, No. 35).




Global changes in falciparum malaria endemicity

The present distribution of *falciparum* malaria is captured well (78% accuracy) by five climate variables, minimum and maximum temperature conditions, precipitation and vapour pressure (Figure 3.4a). Only in Iran and South-East Brazil are there significant areas of false positive predictions (SE Brazil is in fact shown as a low-risk area on some malaria maps). Under the medium-high climate-change scenario (Figures 3.4b-d), the most significant threat to UK citizens travelling abroad is the progressive spread northwards of potentially malarious areas through Mexico into the Southern states of USA. The potential spread of malaria in Florida is particularly serious given the tourist industry and the common perception of zero-threat there, although the Florida's very active anti-mosquito services would undoubtedly react swiftly. Similarly, South-west Turkey, an increasingly important holiday destination, could become a high risk region for *falciparum* malaria. Globally, only central Brazil and Venezuela appear to lose their suitability for malaria.


Figure 3.4 Global *falciparum* malaria.

The present-day distribution of *falciparum* malaria is adequately described by temperature, precipitation and vapour pressure data (Fig. 4a, 78% correct predictions, 14% false positives and 8% false negatives). The predicted probabilities with which local climates match those in malarious areas are colour coded according to the inset probability scale. Predictions for the HadCM2 medium-high scenarios of 2020, 2050 and 2080 are shown in Fig4 b), c) and d).


Global malaria. Predicted areas of suitability, 1961-90 climate




Global malaria. Predicted areas of suitability, HadCM2 2020s medium-high scenario



Global malaria. Predicted areas of suitability, HadCM2 2050s medium-high scenario



Global malaria. Predicted areas of suitability, HadCM2 2080s medium-high scenario




These same changes, apart from those in South America, are predicted by the 2050s under all except the lowest scenarios of climate change (Figure 3.5).


Figure 3.5 Global *falciparum* malaria.

As for Figure 3.4, for the Low, Medium-Low, Medium-High and High HadCM2 2050s scenarios.


Global malaria. Predicted areas of suitability, HadCM2 2050s low scenario




Global malaria. Predicted areas of suitability, HadCM2 2050s medium-low scenario



Global malaria. Predicted areas of suitability, HadCM2 2050s medium-high scenario



Global malaria. Predicted areas of suitability, HadCM2 2050s high scenario



- Certain parts of the world frequently visited by tourists from the UK, especially those on package tours, will become increasingly suitable for *falciparum* malaria in the future. These areas include Florida and other Southern states of the USA and South-West Turkey. UK health authorities should monitor the variable success of the local health authorities at these holiday destinations, to combat any increasing malaria risk to UK travellers abroad.

3.4 Tick-Borne Infections

Tick-borne pathogens are almost all zoonoses, circulating naturally amongst wild vertebrate hosts, but also infecting humans that are accidentally bitten by the vector tick. In Europe, the tick *Ixodes ricinus* is the principal vector of two pathogens that frequently infect humans: bacteria of the *Borrelia burgdorferi* complex, that cause Lyme disease, and the virus that causes tick-borne encephalitis (TBE). In addition, humans are occasionally infected with viruses that cause mortality in sheep, e.g. louping ill virus, and with protozoa, *Babesia* spp., that cause redwater fever in cattle.

The major biological risk factors for all tick-borne infections are the distribution, abundance and pattern of seasonal activity of the vector ticks (Randolph, 1998). Ticks feed only once per life stage, as a larva, a nymph and an adult, between which they spend long developmental and host-seeking periods on the ground. Not all hungry ticks are infected. Nymphs are regarded as the most significant risk, because infection prevalence in them is usually higher than in larvae, having been amplified as larvae feed on infected hosts. Furthermore, nymphs are more abundant and smaller (less noticeable) than adults.

Interacting with these risk factors are human activities, outdoor occupations and leisure pursuits that bring humans into contact with ticks.

Lyme disease (predicted using the expert opinion method)

Lyme disease is widespread and prevalent throughout Europe and the UK, occurring more or less wherever ticks occur. The number of cases reported annually in UK, <200, (O'Connell *et al*, 1998) is far lower than in mainland Europe despite similar densities of infected ticks. One explanation is that a large number of cases still go undetected and unreported, although UK GPs are increasingly aware of the problem. There is some evidence that the strain of *Borrelia* in UK is different: intensity of infection in ticks is very low and isolation of live bacteria is very much more difficult (Livesley *et al*, 1994). Perhaps this results in lower rates of transmission to humans. Furthermore, the infection prevalence in ticks is geographically highly variable, depending on host factors (see below).

A climate-induced change in risk in the UK?

There is no simple correlation between temperature and incidence of Lyme disease in UK. Data presented in Cannell *et al* (1999) show that the annual number of cases reported in UK from 1986 to 1997 have increased since 1994, but there is no significant correlation with mean summer temperatures in Central England.

Tick distribution

Ticks are currently distributed throughout the UK, from Northern Scotland to the South coast. Climatically, therefore, the whole of the UK is suitable for ticks. Their presence in

any one locality is determined principally by the presence of suitable hosts for the adult ticks; while larvae and nymphs feed on vertebrates of all sizes, from mice and blackbirds to pheasants and deer, adults are confined to large mammals such as deer, sheep and cattle. In addition, the habitat structure must afford the right micro-environment for tick survival, with good vegetation cover above a substantial litter or mat layer that retains moisture. This usually coincides with deer habitats (deciduous woodlands) (Daniel *et al*, 1998) and rough grazing. Improved pastures for sheep grazing are rendered unsuitable for ticks.

Any increase in tick-infested areas is more likely to be the effect of changing agricultural and wildlife management practices than the effect of changes in climate alone. This has been seen in Scotland, for example, where grouse moors have been invaded by bracken. Increasing deer populations will support ticks in more places (as appears to have happened in the North-East USA).

Tick abundance

Ticks are most abundant where:

- a) hosts, particularly for the adult stage, are abundant;
- b) the overall climate is warm enough to allow rapid development between tick stages and sufficiently wet to allow good survival between feeds;
- c) the vegetation creates a favourable microclimate.

- It is almost impossible to predict reliably any change in tick abundance with climate change, because it will be the outcome of two opposing forces: higher temperatures, especially overwinter, will accelerate development and may eliminate diapause (a period of winter quiescence, when ticks do not feed), while drier summers will limit tick host-seeking activity and increase mortality directly. Only a climate-driven population model, which is not yet available for *I. ricinus*, will answer this question.

Seasonal patterns of tick host-seeking activity

In warmer regions ticks quest for hosts over longer periods of the year, starting earlier in the spring and continuing later into the autumn. During the summer, unusually dry spells cause a sharp decline in questing activity, from which the ticks may not recover if the dry spell is prolonged.

- It is possible that: the main tick activity season may shift to earlier in the spring; there may be a more pronounced autumn peak; and numbers of questing ticks may decline more dramatically in mid-summer.

Host factors

Infection prevalence in ticks depends on the specific tick-host relationships in any locality, which determines the circulation of the genetically diverse *Borrelia burgdorferi* complex. The four genotypes found in the UK have distinct transmission cycles *via* different vertebrate hosts (Kurtenbach *et al*, 1998). Pheasants, for example, are only competent to transmit two genotypes, *B. garinii* associated with neurological disorders (e.g. Bell's palsy) and *B. valaisiana* that has no associated pathology. In woodlands with large populations of pheasants, circulation of the other genotypes *via* mammals appears to be inhibited. Moreover, because pheasants feed mostly nymphal ticks, the prevalence of infection is high only in the relatively few adult ticks which emerge from these fed nymphs. In woodlands where rodents and squirrels feed most immature tick stages, nymphs have a much higher prevalence of infection with *B. afzelii* (associated with cutaneous symptoms) and *B. burgdorferi* s.s. (associated with arthritis) (Craine *et al*, 1997; Humair and Gern, 1998). On moorlands dominated by sheep, the specific transmission cycle again results in high infection prevalence only in adult ticks (Ogden *et al*, 1997).

- Unless the predicted climate change has an impact on vertebrate fauna (which is possible) the important risk factor of the type and abundance of the various Lyme disease vertebrate hosts will not be affected.

Human behaviour/activity

With warmer weather, humans are likely to interact with ticks in their habitat more frequently and for longer in the year. The creation of new leisure parks in forested areas should be a cause for some concern, since this will increase the contact rates between humans and ticks.

- People engaging in leisure activities are most likely to contact Lyme disease-infected ticks during spring and autumn, when ticks are likely to be most active. Whether, under various scenarios for climate change, there will be an increase in numbers of infected ticks at these times is still uncertain.

Tick-borne encephalitis (TBE) (predicted by the statistical approach)

At present, TBE is confined to recognisable foci within Central Europe, the Baltic region and extensively through the Russian Federation (ImmunoAG, 1997; Korenburg, 1994). Currently there is a real risk of infection to holidaymakers walking in these countries. We now understand that virus circulation depends on a particular pattern of tick seasonal activity (Randolph *et al*, 1999) which occurs only in certain parts of the tick's geographical range, where the seasonal temperature profile is typically continental (as opposed to oceanic): high summer temperatures followed by rapid cooling in the autumn (Randolph *et al*, 2000). In regions that are too dry, however, poor tick survival limits TBE virus maintenance.


The extent, although not the focality, of the present distribution can be predicted very well (86% accuracy) from five climatic variables, including four variables of minimum and maximum temperature conditions, together with the maximum vapour pressure as a measure of terrestrial moisture conditions (Figure 3.6a).

A reduced risk of TBE in the future?


The predicted rise in temperature and decrease in moisture in the summer appears to drive the distribution of TBE virus into higher latitude and higher altitude regions progressively through the 2020s, 2050s and 2080s (Figure 3.6b-d). However, the altitude of the alpine region is too great to allow TBE virus to become established there. In the 2020s, France, Switzerland, Slovenia, Hungary and much of Austria are cleared of TBE virus, and the range of this virus (though not necessarily its vector) has contracted to inland regions of the Baltic states. By the 2050s, TBE has moved into areas at present free of infection, notably the mountains on the Slovak/Polish border and further North-west in Scandinavia. Central Europe is virtually cleared of TBE. This is consistent with the conclusion that increased temperatures have already extended the northern limit of *I. ricinus* in Sweden (Lindgren *et al*, 2000). The final toe-hold in the 2080s is confined to a small part of Scandinavia, including new foci in Southern Finland.

Figure 3.6 Tick-borne encephalitis (TBE) in Europe.


The present-day distribution of TBE in Europe is adequately described by temperature and vapour pressure data (Fig. 6a, 86% correct predictions, 12% false positives and 2% false negatives). Predictions for the HadCM2 medium-high scenarios of 2020s, 2050s and 2080s are shown in Fig 3.6 b), c) and d)



TBE in Europe. Predicted areas of suitability, HadCM2 2050s medium-high scenario



TBE in Europe. Predicted areas of suitability, HadCM2 2080s medium-high scenario




For a given date, e.g. 2050, a similar pattern of TBE distribution emerges as different scenarios are considered (Figure 3.7): the TBE virus is pushed to the North-East of its present range, only moving westwards in Southern Scandinavia. Only under the low and medium-low scenarios does TBE remain in central and Eastern Europe to any extent.


Figure 3.7 Tick-borne encephalitis (TBE) in Europe.

As for Fig. 6, for the Low, Medium-Low, Medium-High and High HadCM2 2050s scenarios.


TBE in Europe. Predicted areas of suitability, HadCM2 2050s low scenario




TBE in Europe. Predicted areas of suitability, HadCM2 2050s medium-low scenario



TBE in Europe. Predicted areas of suitability, HadCM2 2050s medium-high scenario



TBE in Europe. Predicted areas of suitability, HadCM2 2050s high scenario



None of the predicted climatic changes suggest that the UK will be threatened by the TBE virus. Overall, there may be a marked decrease in the extent of this pathogen, although some areas free at present may be invaded, notably the highlands of the Czech and Slovak Republics and parts of Scandinavia, north and west of the present coastal endemic regions.

3.5 Possible sources of other vector-borne diseases (predicted by the climate envelope approach)

At present, on the basis of co-varying mean temperature, precipitation and vapour pressure, similar climatic conditions to those found in various parts of the UK are found extensively in Northern Europe, Asia west of the Caspian Sea, Japan and neighbouring regions of China and New Zealand (Figure 3.8a).

Under the medium-high scenario, the changed climate in the UK is predicted to match the present climate in different parts of the world (Figures 3.8b-d). For example, by the 2020s parts of the UK match more of the Mediterranean rim: does this pose a threat of leishmaniasis? Farther into the future for UK climates, the regions of similarity with present-day Europe then decrease; by the 2080s the climate predicted for much of Southern, Central and North-West England does not match any found in Europe today. If climate really is a determining factor for the arrival of new diseases into the UK, this exercise highlights the regions of the world from which we might expect to import problems. With data on which diseases occur in these places at the present time, we should be able to make more informed guesses of the future risks of exotic diseases to inhabitants of the UK.

Figure 3.8a. Matching the UK to Global climates

a) The UK inset shows the distribution of 10 climatic zones defined in terms of temperature, rainfall and vapour pressure (the mean, maxima and minima) for the period 1961-90. The world map shows the global distribution of these same zones in the same period, with the detail for Europe.




Figure 3.8b. Matching the UK to Global climates

b) The UK inset shows the distribution of 10 climatic zones defined in terms of temperature, rainfall and vapour pressure (the mean, maxima and minima) for the HadCM2 2020 medium-high scenario. The world map shows the global distribution of these same zones at the present time (1961-90), with the detail for Europe.




Figure 3.8c. Matching the UK to Global climates

c) As for Figure 3.8b but using the HadCM2 2050 medium-high scenario.





Figure 3.8d Matching the UK to Global climates

d) As for Figure 8b but using the HadCM2 2080 medium-high scenario.



The same exercise has been done for the various scenarios in the 2050s (Figure 3.9a-d).

Figure 3.9. Matching the UK to Global climates

a) As for Figure 3.8b but using the HadCM2 2050 low scenario.




Fig. 3.9. Matching the UK to Global climates

b) As for Figure 3.8b but using the HadCM2 2050 medium-low scenario.




Figure 3.9. Matching UK to Global climates

c) As for Figure 3.8b but using the HadCM2 2050 medium-high scenario.





Figure 3.9. Matching the UK to Global climates

d) As for Figure 3.8b but using the HadCM2 2050 high scenario.



3.5 Other insect-related problems (predicted by the expert opinion method)

We can only list a range of other insect-related problems which may change with the predicted climate changes in the UK.

Of possible high significance:

- *Flies and diarrhoeal diseases - the “buffet factor”.* Contaminative spread of bacteria by nuisance flies is likely to increase with any increase in the abundance of such insects (see also Chapter 4 on foodborne disease).
- *Midges – relevant to tourism.* The midge menace is already well known to local people and holiday makers in Scotland and elsewhere. An extension of warm summer conditions may well increase the seasonal extent and abundance of these insects.
- *Fleas – nuisance factor associated with cats and dogs.* Fleas on domestic animals thrive in warm conditions, such as those associated with domestic central heating; hungry cat fleas are particularly likely to bite humans. An increasingly warm climate will increase such problems.
- *Stinging/biting insects such as bees, wasps and horseflies - serious allergic reactions.* Allergic reactions to bee and wasp stings can be fatal in a minority of cases. Sensitivity appears to increase with increased exposure, a possible outcome of warmer conditions.
- *Urticaceous caterpillars - allergies.* Caterpillars of certain moths occur in vast numbers and live communally, often spinning characteristic silk 'tents' that attract human attention. On contact with human skin, the hairs of such insects can cause painful swellings and rashes that persist for many days. We are uncertain of the causes of outbreaks of such caterpillars, but they may increase in frequency in warmer conditions.

Others to be considered:

- *Nuisance mosquitoes, or migrant pests such as locusts.*
- *Introduced new vector species (e.g. Stegomyia albopicta (Aedes albopictus).* Potential vector of both dengue fever and malaria, accidentally introduced to the USA in used car tyres and spreading rapidly).
- *Black flies (e.g. the Blandford fly) - periodic and newsworthy outbreaks of these blood-sucking flies occur in Southern UK counties.*
- *House dust mites – associated with allergies and possibly with asthma.*
- *Plague – endemic in the USA, transmitted by fleas between rats and also directly from person to person.*
- *Leptospirosis (e.g. Weill’s disease) – already present in the UK, related to direct contact with rodents or with areas contaminated with rodent urine and faeces.*

3.6 Priorities for future action - to identify the most immediate dangers from vector-borne diseases

Since we are dealing with considerable uncertainties concerning vector-borne diseases, we feel there are two priorities for future action. The first is to monitor changing risks as they are happening, both within the UK and to holiday makers. The second is to examine the impact of multivariable environments on the spatial and temporal patterns of disease distribution and intensity for both temperate and tropical vector-borne disease systems.

- Create a database against which to monitor change in incidence.
- GPs to report centrally on any insect-associated conditions, e.g. wasp/bee stings, rashes from caterpillars, tick bites.
- Any significant change in incidence should be the alert for focused research.
- Map the distribution and abundance of key vector species in the UK.

Identify the threat to travellers/holidaymakers visiting the tropics:

- Match holiday destinations to disease risk.
- Monitor risk in those places.
- Alert GPs in the UK to recognise disease symptoms and to report centrally.

Build on studies to date that have used remotely-sensed satellite data for extensive studies of environmental variables that determine vector-borne disease risk (e.g. for malaria, dengue, leishmaniasis, trypanosomiasis and tick-borne encephalitis).

- Identify the regions of high risk that might threaten inhabitants of the UK.
- Conduct satellite data analysis and related field studies to refine the predictions of risk to the UK.

References

- Boyd, M. F. (1949) *Malariology. A comprehensive survey of all aspects of this group of diseases from a global standpoint*. Philadelphia and London: Saunders, W.B
- Cannell, M.G.R., Palutikof, J.P. and Sparks, T.H. (1999) *Indicators of Climate Change in the UK*. Department of the Environment, Transport and the Regions.
- Craine, N.G., Nuttall, P.A., Marriott, A.C. and Randolph, S.E. (1997) Rôle of grey squirrels and pheasants in the transmission of *Borrelia burgdorferi sensu lato*, the Lyme disease spirochaete, in the U.K. *Folia Parasitologica* **44**, 155-160.
- Curtis, C. (2003) Susceptibility of *Anopheles plumbeus* to *Plasmodium falciparum*. Abstract. In *International Congress of Parasitology, 4-6 July 2003*,. Iasi, Romania
- Daniel, M., Kolár, J., Zeman, P., Pavelka, K. and Sádlo, J. (1998) Predictive map of *Ixodes ricinus* high-incidence habitats and a tick-borne encephalitis risk assessment using satellite data. *Exp.Appl.Acarol.* **22**, 417-433.
- Danis, M., Mouchet, M., Giacomini, T., Guillet, P., F, L. and M, B. (1999) Indigenous, introduced and airport malaria in Europe. *Medecine et Maladies Infectieuses* **26**, 393-96.
- Davidson, N., d'A Laffoley, D., Doody, J., Way, L., Gordon, J., Key, R., Pienkowski, M., Mitchell, R., and Duff, K. (1991) *Nature conservation and estuaries in Great Britain*. Peterborough: Nature Conservancy Council.
- Detinova, T. (1962) *Age-grouping methods in Diptera of medical importance*. Geneva: World Health Organisation.
- Dobson, M. J. (1997) *Contours of death and disease in early modern England*. Cambridge: Cambridge University Press.
- Gratz, N., Steffen, R. and Cocksedge, W. (2000) Why aircraft disinfection? *Bulletin of the World Health Organisation* **78**, 995-1004.
- HPA. (2004) Malaria imported into the UK, 2003: implications for those advising travellers: Health Protection Agency.
- Humair, P.-F. and Gern, L. (1998) Relationship between *Borrelia burgdorferi sensu lato* species, red squirrels (*Sciurus vulgaris*) and *Ixodes ricinus* in enzootic areas in Switzerland. *Acta Tropica* **69**, 213-227.
- Hutchinson, R., Bayoh, M. and Lindsay, S. (2005) Risk of airport malaria in the UK. . *European Mosquito Bulletin* **19**, 12-13.
- Immuno A.G. Tick-borne encephalitis (TBE) and its immunoprophylaxis. Vienna, Austria (1997).
- James, S. (1931). Some general results of a study of induced malaria in England. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **24**, 477-525.
- James, S. P. (1920) Malaria at home and abroad. *John Bale, Sons and Danielsson, Ltd. London* Chapter V, 79-93.
- Jetten, T. and Takken, W. (1994) *Anophelism without malaria in Europe. A review of the ecology and distribution of the genus Anopheles in Europe*. Wageningen: Wageningen Agricultural University.
- Karch, S. (1996) Breeding of *Anopheles plumbeus* in tires in France. *Journal of Vector Ecology* **21**, 201-201.
- Korenberg, E.I. (1994) Comparative ecology and epidemiology of Lyme disease and tick-borne encephalitis in the former Soviet Union. *Parasitology Today* **10**, 157-10.
- Kurtenbach, K., Peacey, M.F., Rijpkema, S.G.T., Hoodless, A.N., Nuttall, P.A. and Randolph, S.E. (1998) Differential transmission of the genospecies of *Borrelia burgdorferi sensu lato* by game birds and small rodents in England. *Applied and Environmental Microbiology* **64**, 1169-1174.
- Lindgren, E., Talleklint, L. and Polfeldt, T. (2000) Impact of climatic change on the northern latitude limit and population density of the disease-transmitting European tick, *Ixodes ricinus*. *Environmental Health Perspectives* **108**:2;119-123.
- Lindsay, S.W. and Birley, M.H. (1996) Climate change and malaria transmission. *Annals of Tropical Medicine and Parasitology* **90**, 573-588.
- Linton, Y.-M., Lee, A. and Curtis, F. (2005) Discovery of a third member of the *Maculipennis* complex in SW England. *European Mosquito Bulletin* **19**, 5-9.

- Livesley, M.A., Carey, D., Gern, L. and Nuttall, P.A. (1994) Problems of isolating *Borrelia burgdorferi* from ticks collected in United Kingdom foci of Lyme disease. *Medical and Veterinary Entomology* **8**, 172-178.
- Newman, G. (1919) *Reports and papers on malaria contracted in England, 1918*. (12): Local Government Board Public Health Reports.
- Nuttall, G.H., Cobbett, L. and Strangeways-Pigg, T. (1901) The geographical distribution of anopheles in relation to the former distribution of ague in England. *J. Hygiene, Cambridge*. 1, 4-44.
- O'Connell, S., Granström, M., Gray, J.S. and Stanek, G. (1998) Epidemiology of Lyme borreliosis. *Zentralblatt für Bakteriologie* **287**, 229-240.
- Ogden, N.H., Nuttall, P.A. and Randolph, S.E. (1997) Natural Lyme disease cycles maintained *via* sheep by co-feeding ticks. *Parasitology* **115**, 591-599.
- Ramsdale, C.D. and Coluzzi, M. (1975) Studies on the infectivity of tropical African strains of *Plasmodium falciparum* to some southern European vectors of malaria. *Parassitologia* **17**, 39-48.
- Randolph, S.E. (1998) Ticks are not insects: consequences of contrasting vector biology for transmission potential. *Parasitology Today* **14**, 186-192.
- Randolph, S.E. (2000) Ticks and tick-borne disease systems in space and from space. *Advances in Parasitology* **47**, 217-243.
- Randolph, S.E., Miklisová, D., Lysy, J., Rogers, D.J. and Labuda, M. (1999) Incidence from coincidence: patterns of tick infestations on rodents facilitate transmission of tick-borne encephalitis virus. *Parasitology* **118**, 177-186.
- Rogers, D.J., Hay, S.I. and Packer, M.J. (1996) Predicting the distribution of tsetse flies in West Africa using temporal Fourier processed meteorological satellite data. *Annals of Tropical Medicine and Parasitology* **90**, 225-241.
- Shute, P. G. and Maryon, M. (1974) Malaria in England past, present and future. *Royal Society of Health* **1**, 23-29 and 49.
- Simini, B. (1997) First case of indigenous malaria reported in Italy for 40 years. *Lancet* **350**, 717.
- Snow, R., Craig, M., Newton, C. and Steketee, R. (2003) The public health burden of *Plasmodium falciparum* malaria in Africa: Deriving the number. Working Paper No. 11, Disease Control Priorities Project. Bethesda, Maryland: Fogarty International Center, National Institutes of Health. August 2003.
- Snow, R., Guerra, C., Noor, A., Myint, H. and Hay, S. (2005) The global distribution of clinical episodes of *Plasmodium falciparum* malaria. *Nature* **434**, 214 – 217.
- Targett, G., Drakeley, C., Jawara, M., von Seidlein, L., Coleman, R., Deen, J., Pinder, M., Doherty, T., Sutherland, C., Walraven, G. and Milligan, P. (2001) Artesunate reduces but does not prevent post treatment transmission of *Plasmodium falciparum* to *Anopheles gambiae*. *Journal of Infectious Diseases* **183**, 1254-1259.
- Williams, J., Chitre, M. and Sharland, M. (2002) Increasing *Plasmodium falciparum* malaria in southwest London: a 25 year observational study. *Archives of Disease in Childhood* **86**, 428-430.

Annex.

The following text explains the derivation of the maps in Fig. 1, using a biological approach.

Predicting the areas of suitability for European *vivax* malaria using a biological approach

The analyses to produce the maps of European *vivax* malaria in Figure 3.1 are based on the concept of the basic reproduction rate (R_0), which represents the number of future cases of malaria derived from one infective case at the present time, before this case is cured, or the infected person dies. Where R_0 is greater or equal to 1.0 the disease can become established; when it is less than 1.0 it eventually becomes extinct. One expression for R_0 is shown below:

$$R_0 = \frac{ma^2bp^n}{-\ln(p)r}$$

where,

ma = the number of bites per person per day/night. This was set equal to 1.0 in the present model, because it was assumed that the maximum biting rate tolerated by people living near these areas would be one bite each night, since we reasoned that most people would avoid being bitten more than 30 times in a month.

a = the frequency of feeding on a person, expressed as a daily rate;

$$a = \frac{h}{u} \text{ bites/person/day}$$

where ***h*** is the proportion of mosquito blood meals taken from people (as opposed to other animals that are not infected with human malaria) and ***u*** is the length in days of the gonotrophic cycle - the interval between each egg-batch and, generally, each mosquito blood meal. The present model assumes a mean value of ***h*** of 0.42 for indoor-resting mosquitoes (e.g. *An maculipennis*, in Jetten and Takken, 1994). ***u*** is length of the gonotrophic cycle, described as follows:

$$u = \frac{f_1}{T - g_1} \text{ days}$$

Where ***f*₁** is a thermal sum, measured in degree days, representing the accumulation of temperature units over time to complete the cycle = 36.5°C, ***g*₁** is a development threshold below which development ceases = 9.9°C, and ***T*** is ambient temperature (Detinova, 1962).

p = the daily survival probability of adult mosquitoes. The present model takes the median value of the mortality rate for *An. atroparvus* = 0.029/day (n = 24, range 0-0.294/day) (Jetten and Takken, 1994).

n = the period of parasite development within the adult mosquitoes, in days (the sporogonic cycle).

$$n = \frac{f_2}{T - g_2} \text{ days}$$

Where ***f*₂** is a thermal sum, measured in degree days, representing the accumulation of temperature units over time to complete the development = 105 degree days, ***g*₂** is a development threshold below which development ceases = 14.5°C and ***T*** is ambient temperature (Jetten and Takken, 1994).

b = the proportion of vector females developing parasites after taking an infective blood meal. The model assumed a value of 0.19 (James 1951).

r = the rate of recovery of humans from infection with malaria. The usual assumption is that the duration of each infection is therefore $1/r$ days. The model assumed that an infection would be patent for 60 days, giving a value for **r** of 0.0167/day (Boyd, 1949).

In the model the above formulae were used together with the various scenarios for climate change in the UK. The model output the number of months of the year when **R₀** is greater than 1.0, indicating potential disease spread. Under conditions when **R₀** is less than 1.0 for a considerable proportion of the year, the disease probably cannot persist without continuous introduction from elsewhere, or possibly as quiescent stages within apparently recovered people.

4 Foodborne disease and climate change

(See also section 4.2 of 2001 report)

Graham Bentham, School of Environmental Sciences, University of East Anglia

Summary

- New scientific evidence confirms the effect of temperature on salmonellosis.
- The role of temperature in *Campylobacter* transmission remains uncertain.
- Effect of warmer summers on food borne disease incidence will depend on future food hygiene behaviour, the relative contribution of different pathogens, as well as changes in temperature.

Recommendations

- Vulnerability to the effects of climate change could be reduced by continuing efforts to improve the microbiological standards of food at all stages in the food chain including production, distribution, storage and preparation.
- There may be a case for educating and warning food producers and the public about the particular risks associated with hot weather.
- The existence of lags between high temperatures and effects on humans offer the possibility of instituting a warning system.

4.1 Introduction

The previous review noted that food poisoning is an important cause of morbidity in the general population associated with significant costs of treatment and loss of working time. The 1990s had seen a large increase in incidence for reasons that remain controversial but may include changes in methods of food production such as the shift towards intensive rearing of poultry and other animals, changing patterns of retailing and catering as well as changing behaviour of consumers.

The recognition of the seasonality of food poisoning incidence and of the various ways in which weather conditions might affect the micro-biological safety of food had led to the suggestion that climate change could increase food poisoning risk (McMichael *et al*, 1996). However, at the time of the previous review there was only a very limited body of published evidence on which an assessment of the potential impacts of climate change on food poisoning could be based.

In recent years there have been major changes in the number of cases of food poisoning in England and Wales and in the relative importance of different pathogens. The number of food poisoning notifications increased steadily during most of the 1990s to reach a peak in 1998. Since then there has been a marked decline with notifications for the most recent year (2004) being about 25% lower than at the peak with especially large reductions being evident for infections from *Salmonella* which decreased by 60% between their peak (in 1997) and 2004.

4.2 Current impacts of climate and weather


The existence of a pronounced seasonal pattern, with a higher incidence in the summer, points to a role for weather and climate in the aetiology of food poisoning. High temperatures favour the multiplication of some pathogenic micro-organisms in food including the *Salmonellas* that are an important source of food poisoning in the United Kingdom. High temperatures may also have an influence on human health risks by affecting infection rates in food animals, for example by the multiplication of bacteria in animal feed. Other indirect influences on human risks could include a weather influenced shift towards dietary items or forms of food preparation (e.g. barbecues) that are associated with increased risk.

Since the previous review, several empirical studies have been published on the relationship between environmental temperatures and the incidence of foodborne illness. Bentham and Langford (2001) analysed the association between weekly notification of food poisoning and temperatures in England and Wales for the period 1974-96. This confirmed the findings of earlier studies (Bentham and Langford, 1995; Bentham, 1997) of a positive association between food poisoning and temperature with the strongest effects being for conditions 2-5 weeks earlier. A significant limitation of this study is that the data used do not record which pathogens are responsible for the notified cases. This is important because the different common types of food poisoning pathogens are known to respond differently to environmental temperatures. The rate of multiplication of *Salmonella* is strongly related to temperature in the range from approximately 7 to 37°C providing opportunities for ambient temperatures to affect the numbers of bacteria in food at various stages in the food chain. *Campylobacter*, on the other hand, replicates most readily at temperatures between 37 and 42°C in a low oxygen environment. This makes it well adapted to the guts of birds and other animals but multiplication in food at typical ambient temperatures is unlikely. Simple direct relationships with environmental temperatures are therefore less likely than for *Salmonella*, although more complex effects of

temperature on the ecology of animal reservoirs or on patterns of human behaviour leading to exposure are possible.

D'Souza *et al*, (2003) investigated the association between monthly salmonellosis notifications and temperature in 5 Australian cities. They found a significant positive association between mean temperature of the previous month and the number of salmonellosis notifications in the current month with the estimated increases for a 1°C ranging from 4% to 10% depending on the city. Kovats *et al*, (2004) analysed the association between laboratory-confirmed cases of salmonellosis and monthly temperatures in 10 European countries. The estimated change in incidence above a common 6°C threshold ranged from 0.3% in Denmark to 12.5% in England and Wales. The strongest effects were found for temperatures 1 week before the onset of illness rather than the longer lag of 1 month found in the Australian study.

Figure 5.1. Modelled association between temperature and number of reported cases of salmonellosis in England and Wales (adjusted for outbreaks, seasonal factors and holidays). Source: Kovats *et al*, 2004.



Campylobacter

At the time of the earlier review, information was sparse on the relationship between environmental temperatures and human disease resulting from infection by *Campylobacter*, but since then new evidence has been published. In a study of *Campylobacter* infection in Denmark, Patrick *et al*, (2004) found a significant positive association with temperature, the strongest effects being for lags of 4 weeks. The study also showed that the prevalence of *Campylobacter* infection in broiler flocks at slaughter was positively associated with temperatures 3 weeks earlier. In Denmark consumption of poultry is an important risk

factor for campylobacteriosis in humans. Increasing *Campylobacter* carriage rates in chicken meat following warm weather may therefore be one of the factors underlying the association between incidence and temperature in humans. A positive association between rates of campylobacteriosis in humans and temperature (unlagged) has also been reported for England and Wales (Louis *et al*, 2005). However, a limitation of both of these studies is that they did not adjust for seasonal factors that could confound the association between campylobacteriosis and temperature. Influences on the seasonality of campylobacteriosis in 13 countries has been investigated by Kovats *et al*, (2005). Although most of these countries showed seasonal patterns the study was unable to find a strong effect of temperature variability on campylobacter transmission. Unlike *Salmonella* a simple direct relationship between temperature and *Campylobacter* infections is therefore not likely. However, what cannot be excluded is the possibility of more complex effects of weather and climate via their influence on factors such as the size of animal reservoirs of infection (Skelly and Weinstein, 2003) or the ease of transmission to humans for example by vectors such as flies (Nicholls, 2005).

4.3 Future impacts of climate change

One study (Bentham, 1997) had used regression analysis to develop statistical models of the relationship between the monthly incidence of food poisoning and temperatures in England and Wales for the period 1975-95. The resulting regression model of the association between temperature and food poisoning notifications (adjusted for trend and seasonal factors) was then used to estimate the impact on food poisoning notifications of scenarios of +1, +2 and +3°C temperature increases. This produced estimates of increases in food poisoning notifications of 4.5%, 9.5% and 14.8% respectively. If applied to the total of 94,000 notified cases of food poisoning for 1998 these would represent absolute increases of ~4000, 9000 and ~14,000. It was emphasised that because of the under-recording of food poisoning the real number of additional cases might be considerably higher. It was concluded that higher temperatures as a result of climate change might exacerbate the food poisoning problem which is already a significant threat to public health. However, it was emphasised that the estimated changes were small relative to the large increases that had occurred over the previous twenty years as a result of other factors.

The 2001 report concluded that a 1°C increase in temperature might result (*ceteris paribus*) in about a 4.5% increase in food poisoning. Subsequent published studies have strengthened the evidence that environmental temperatures affect the risk of food poisoning by *Salmonella* with Kovats *et al*, (2004) yielding an estimated effect of +12.5% per 1°C increase in temperature. This could be taken to indicate that the earlier estimate is too low. However, the impacts on total incidence of food poisoning resulting from all pathogens will clearly depend also on the temperature sensitivity of disease caused by other pathogens, especially *Campylobacter*. Unfortunately, in spite of the new data that have become available the published evidence remains insufficient to reliably estimate the potential effect of changes in temperature on campylobacteriosis. Another important factor is that in recent years both the absolute and the relative contribution of *Salmonella* infections to the total burden of food poisoning have declined. The fraction of food poisoning cases for which there is clear evidence of a direct relationship with temperature is therefore likely to have fallen. On balance, there appear to be no strong grounds for changing the estimate of the earlier review that a 1°C increase in temperature might result in about a 4.5% increase in food poisoning. The absolute numbers of additional cases will depend on the scenarios for temperature and the baseline number of cases of food poisoning.

References

- Bentham, G. and Langford, I.H. (2001) Environmental temperatures and the incidence of food poisoning in England and Wales. *Int.J.Biometeorol.* **45**,22-6.
- D'Souza, R.M., Becker, N.G., Hall, G. and Moodie, K.B.A. (2003) Does ambient temperature affect foodborne disease? *Epidemiology* **15**;1:86-92.
- Kovats, R.S., Edwards, S.J., Hajat, S., Armstrong, B.G., Ebi, K.L. and Menne, B. (2004) The effect of temperature on food poisoning: a time-series analysis of salmonellosis in ten European countries. *Epidemiol.Infect.* **132**:443-53.
- Kovats, R.S., Charron, D., Cowden, J., D'Souza, R.M., Ebi, K.L., Edwards, S.J., Gauci, C., Gerner-Smidt, P., Hajat, S., Hales, S., Hernández Pezzi, G., Kriz, B., Kutsar, K., McKeown, P., Mellou, K., Menne, B., O'Brien, S., van Pelt, W. and Schmid, H. (2005) Climate variability and campylobacter infections: an international study. *International Journal of Biometeorology*, **49**(4):207-214.
- Nichols, G.L. (2005) Fly transmission of Campylobacter. *Emerg.Infect.Dis.* **11**;3:361-4.
- Patrick, M.E., Christiansen, L.E., Wainø, M., Ethelberg, S., Madsen, H. and Wegener, H.C. (2004) Effects of climate on incidence of *Campylobacter* spp. in humans and prevalence in broiler flocks in Denmark. *Applied and Environmental Microbiology* **70**:7474-80.
- Skelly, C. and Weinstein, P. (2003) Pathogen survival trajectories: an eco-environmental approach to the modelling of human campylobacteriosis ecology. *Environmental Health Perspectives* **111**;1:19-28.

5 Water and disease and climate change

(See also section 4.4 of 2001 report)

Gordon Nichols, Centre for Infections, Health Protection Agency
Sari Kovats, London School of Hygiene and Tropical Medicine

Summary

- Climate change is likely to impact on raw water quality, which in turn could affect treatment efficiency and the stability of drinking water in distribution.
- More frequent extreme rainfall events could lead to increased surface water turbidity, and higher numbers of indicator bacteria and pathogens in surface water. This would create a greater challenge for water treatment works particularly where direct river abstraction is used.
- Changes in the distribution of rainfall may cause local increases in drought. In such areas, there may be an increased likelihood of contaminated surface water reaching groundwater through the opening of short circuits as a result of a lowered water table.
- Increased temperature and nutrient concentrations in surface waters could lead to cyanobacterial blooms occurring more often.
- The greatest impact of increased temperature would appear to be on the efficiency of chemical coagulation, which may lead to decreased removal of microbes by subsequent clarification and filtration. The result could be an increased challenge on final disinfection.
- These potential impacts of climate change are unlikely to pose a threat to well managed water treatment plants, but could cause problems for private water supplies, surface water supplies without filtration and groundwater supplies under the influence of surface water, unless they are adequately filtered.
- Changes in surface and ground water availability must be carefully managed to avoid acute crisis in provision at the household level.

Recommendations

- Improved risk management of water resources in the UK to avoid acute crisis in supply.
- Risk assessment by water companies of extreme weather events, particularly intense rainfall.
- Integrated monitoring of algal blooms in surface waters.
- Control of filter operation to minimise breakthrough of filters should be improved
- Monitoring filter performance by means of turbidimeters should be increased.

5.1 Introduction

The potential impact of climate change on waterborne infectious diseases has not been particularly well addressed in climate change assessments, in contrast to the treatment of vector-borne diseases. The conclusions of the 2001 report were limited by a lack of research of the effect of weather and climate on waterborne disease in the UK. Since then, the water industry and the Environment Agency have taken steps to assess potential risks associated with climate change (Nichols *et al*, 2003; Wilby *et al*, 2005).

5.2 The effects of heavy rainfall on outbreaks of water borne disease

Microbiological contamination of public supplies is an important concern in Europe. However, few studies have investigated an association between rainfall and human disease. Due to the complexity of the causal pathways on the route from surface or groundwater to the household, it is difficult to detect what may be a small contribution to the overall burden of disease.

Several studies have investigated an association between drinking water turbidity and health. Although the methodologies have been criticised, there is some indication that reducing the maximum turbidity of filtered water may reduce the risk of *Cryptosporidium* contamination. Rainfall appears to increase concentrations of *Giardia* and *Cryptosporidium* in surface water through its effect on the amount of solid matter in the water. There may be a link between when *Cryptosporidium* and *Giardia* concentrations peak and when, for instance, turbidity reaches its highest level. Open finished water reservoirs are at risk for post-treatment faecal contamination by animals, including those that shed species of *Giardia* and *Cryptosporidium* that are potentially pathogenic to humans. A study carried out in the US found a statistically significant association between extreme rainfall events and monthly reports of outbreaks (Curriero *et al*, 2001).

Heavy rainfall can cause abnormal changes in the direction of flow of water through both surface and underground channels. For example, in one outbreak of cryptosporidiosis associated with a borehole extracted groundwater, it was thought that the heavy rainfall led to water running across the surface of a fields where cattle were grazing (Bridgman *et al*, 1995). The water (and cattle faeces) then pooled around the head of the borehole and leaked into the water supply.

In impermeable catchments, microbial contaminants on the land will be carried into rivers more rapidly than those present on permeable catchments, particularly following periods of dry weather during which the surface has become “baked”. One of the predicted consequences of climate change will be more severe rainfall events. In these situations, microbial contaminants present in biosolids or manures applied to agricultural land may be transferred to surface water sources more rapidly than under conditions of more “normal” rainfall. The contamination of surface water sources used for drinking water production by storm drain overflow may be brief with a bolus of infected water being followed by substantial dilution as a result of the excess water flow. Well run water utilities should be able to plan for these events and have a system that keeps the risks of contamination low, irrespective of weather.

Cryptosporidiosis is the most significant water-borne disease associated with the public water supply in the UK. The Spring peak in cases of cryptosporidiosis is likely to be associated with spring rainfall, and high levels of contaminated manure on the land, especially from lambs (Lake *et al*, 2005). These relationships show that rainfall may have played an important role in sporadic cases of the disease in the spring time. The

response by the water industry to the UK *Cryptosporidium* Regulations has led to a reduction in cryptosporidiosis and to fewer outbreaks of this disease (Sopwith *et al*, 2005). Although outbreaks of drinking water related disease do still occur, within the last few years they have nearly all been due to *Cryptosporidium* and are associated with supplies that are inadequately treated. It is likely that the Water Safety Plan approach being sponsored by WHO will lead to further improvements to water quality. There are still many private supplies in use in the UK and these are frequently poorly treated. The microbiological quality of such supplies is generally very poor and can deteriorate markedly after heavy rainfall (Rutter *et al*, 2000). A greater number of extreme weather events could worsen this situation although probably not substantially.

5.3 The effect of drought on water quality and water availability

A real concern is the effect of decreases in water availability. Changes in surface and groundwater availability, particularly in the South of England, will need to be carefully managed to avoid any threats to domestic water supplies. Water shortages can, if they are severe, have dramatic effects on populations. The shortages in the supply of drinking water through Yorkshire Water in 1995 highlighted the limitations of planning using historical records in an era of changing climate.

Warmer wetter winters and drier summers will influence river flows. These are one of the main indicators used to monitor the impacts and progress of climate change. Variable river flows are obviously important since over sixty percent of the UK's water supplies are abstracted from rivers. A recent project funded by the Environment Agency assessed the impact of climate change on effective monitoring of chemical standards for health protection (Crane *et al*, 2005). In the near term, climate change is unlikely to directly cause any exceedances of health and environmental standards. However, unless more samples are taken, changes in climate that lead to an increase in the variability of river flows could lead to an increased likelihood of failure to detect non-compliance with the standards.

Flows in rivers draining impermeable catchment areas decline rapidly when there is little rain and effluent discharges become an increasingly high proportion of the river flow. In contrast, where natural groundwater inflows sustain river flows, the effect of low rainfall will not be so immediate. However, once groundwater levels become low the river flow will be depressed and may remain in this state long after rainfall has occurred. If summers become warmer and drier, effluent discharges could remain an important component of the river flow for protracted periods.

In addition to threatening water supplies, low flows reduce effluent dilution. A greater effluent loading in the river will also be reflected in an increased pathogen loading. This could represent an increased challenge to water treatment plants. However under low flow conditions, transit times between discharge and abstraction will be increased providing more time for pathogen decay to occur. In addition, if climate change brings warmer drier summers, river temperatures will be higher with greater rates of natural purification through biological predation and possibly increased levels of sunlight. An assessment of the dry summer of 2003, when river flows were reduced, found apparent changes in water quality (Senhorst and Zwolsman, 2005). There are also risks of surface water reaching groundwater through routes that are not usually open during periods when the water table is low as a result of a drought.

The most significant consequence of drought would be a failure of the domestic water supply, resulting in a need for standpipes and other methods for water delivery. The potential health effects of this would include infectious intestinal diseases, due to contamination of water and reduced hygiene, but such measures may also lead to civil

unrest. Access to sufficient water for the elderly, disabled and less mobile would be a concern.

5.4 Recreational waters and algal blooms

Recreational waters (either inland surface waters or coastal waters) are also an important source of waterborne diseases in bathers (Zmirou *et al*, 2003). Transmission of water borne disease from faecal contamination, as well as naturally occurring pathogens, may also be affected by warmer temperatures as well as changes in rainfall and runoff (Hunter, 2003; Schijven and de Roda Husman, 2005).

Algal blooms may become more common with increasing temperatures. Several species have been implicated in causing disease in humans. However, there is little direct evidence that *A. hydrophila* (or other organisms that bloom in distribution) currently cause any significant burden of ill health. There has not been further work on this since the 2001 report. Some risk assessment is needed, both for direct contact, economic implications (since water bodies must be closed), as well as contaminants in the public water supply (microcystins) (Codd *et al*, 2005). Blooms of *Aeromonas hydrophila* and coliforms can result from raised organic carbon present in the water in distribution, higher temperatures, and reduced chlorine.

There is also evidence that heavy rainfall events can precipitate coliform re-growth in water distribution systems by increasing the nutrient content of water (LeChevallier *et al*, 1991). Non-faecal coliforms in drinking water do not appear to be associated with disease in the community (Edberg *et al*, 1986; Zmirou *et al*, 1987; Hellard *et al*, 2001). It remains possible that other organisms might take advantage of these changed conditions. Public utilities in charge of recreational waters should therefore be vigilant in consulting the organic carbon content of water and should ensure adequate disinfectant levels to prevent these problems.

Cholera is not generally a problem within developed countries. There is a chance that a substantial alteration in world climate could contribute to the triggering of an 8th pandemic of cholera, particularly if there is an increase in sea level. While this might give rise to an increase in the incidence of cholera in people travelling overseas, it is unlikely that indigenous disease would increase with current UK infrastructure. Substantial coastal inundation due to sea level rise might increase population movement and could have a significant impact on immigration and the diseases associated with this.

References

- Bridgman, S.A., Robertson, R.M., Syed, Q., Speed, N., Andrews, N. and Hunter, P.R. (1995) Outbreak of cryptosporidiosis associated with a disinfected groundwater supply. *Epidemiol.Infect.* **115**;3:555-566.
- Codd, G.A., Morrison, L. and Metcalf, J. (2005) Cyanobacterial toxins: risk management for health protection. *Toxicology and Applied Pharmacology* **203**:264-272.
- Crane, M., Whitehouse, P., Comber, S., Ellis, J. and Wilby, R. (2005) Climate change influences on environmental and human health chemical standards. *Human and Ecological Risk Assessment* **11**;2:289-318.
- Curriero, F., Patz, J. A., Rose, J.B. and Lele, S. (2001) The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948-1994. *American Journal of Public Health* **91**;8:1194-1199.
- Edberg, S.C., Piscitelli, V. and Cartter, M. (1986) Phenotypic characteristics of coliform and noncoliform bacteria from a public water supply compared with regional and national clinical species. *Appl.Environ.Microbiol.* **52**;3:474-478.
- Hellard, M.E., Sinclair, M.I., Forbes, A.B. and Fairley, C.K. (2001) A randomized, blinded, controlled trial investigating the gastrointestinal health effects of drinking water quality. *Environ.Health Perspect.***109**;8:773-778.
- Hunter, P.R. (2003) Climate change and waterborne and vectorborne disease. *J Appl.Microbiol.* **94**;s1:37-46.
- Lake, I., Bentham, G., Kovats, R.S. and Nichols, G. (2005) Effects of weather and river flow on cryptosporidiosis. *Water and Health.* **3**;4:469-474.
- LeChevallier, M. W., Schulz, W. and Lee, R. G. (1991) Bacterial nutrients in drinking water. *Appl.Environ.Microbiol.* **57**;3:857-862.
- Nichols,G, Stanfield, G., Hall, T. and Hunter, P.R. Review of the Microbial Implications of Climate Change for the Water Industry. 2003. UK Water Industry Research. Ref Type: Report.
- Rutter, M., Nichols, G.L., Swan, A. and De Louvois, J. (2000) A survey of the microbiological quality of private water supplies in England. *Epidemiol.Infect.***124**;3:417-425.
- Schijven, J.F. and de Roda Husman, A. M. (2005) Effect of climate changes on waterborne disease in the Netherlands. *Water Science and Technology* **51**;5:79-87.
- Senhorst, H.A.J. and Zwolsman, J.J.G. (2005) Climate change and effects on water quality: a first impression. *Water Science and Technology.* **51**;5:53-59.
- Sopwith, W., Osborn, K., Chalmers, R. and Regan, M. (2005) The changing epidemiology of cryptosporidiosis in North West England. *Epidemiol.Infect.* **133**;5:785-793.
- Wilby, R., Hedger, M. and Orr, H.G. (2005) Climate change impacts and adaptation: A science agenda for the Environment Agency of England and Wales. *Weather.* **60**:206-211.
- Zmirou, D., Ferley, J.P. Collin, J.F. Charrel, M. and Berlin, J. (1987) A follow-up study of gastro-intestinal diseases related to bacteriologically substandard drinking water: *Am.J.Public Health.* **77**;5:582-584.
- Zmirou, D., Pena, I., Ledrans, M. and Letertre, A. (2003) Risks associated with the microbiological quality of bodies of fresh and marine water used for recreational purposes: summary estimates based on published epidemiological studies. *Archives of Environmental Health.* **58**;11:703-711.

6 Direct effects of rising temperatures on mortality in the UK

(See also section 4.1 of 2001 report)

GD Donaldson, Queen Mary, University of London

WR Keatinge, Queen Mary, University of London

Summary

- Mean annual heat related mortality did not rise as summers warmed from 1971 to 2003, implying an increase in population tolerance to heat, while annual cold-related mortality fell by more than 33%.
- Daily mortality in the South of England increases at temperatures above about 27°C, and the effect of several consecutive days combined is much greater than the effect of the same number of separate hot days.
- Improved tolerance to heat in the future will reduce the impact of hotter summers, but increased frequency and intensity of heatwaves are a major concern.
- Extrapolation of current data showed the risk of a nine day heatwave (averaging 27°C in South-East England, with over 3000 immediate heat-related deaths and 6350 heat related deaths throughout that summer in Britain), as one in forty each year by 2012, or 25% at some time during the ten years centred on that year.

Recommendations

- Promotion of measures to avoid heat stress and dehydration, on the lines of the current Heatwave Plan, are recommended to reduce heatwave mortality over the next decade.
- In the longer term, improved building design could minimise need for increased energy usage, and consequent acceleration of climate warming by air conditioning.
- Regular assessments of heatwave impacts will be important as climate warming proceeds.

6.1 Introduction

Heat related mortalities are substantial throughout Europe, but the hot summers in Southern Europe cause little more mortality than the milder summers of more northerly regions (Keatinge *et al*, 2000). People in Southern Europe achieve this largely by simple traditional measures, such as use of shade, fans, and avoidance of exercise at hot times of day, rather than by air conditioning as in the United States (Greenburg *et al*, 1983; Rogot *et al*, 1992; Taylor and Goodwin, 2000; Basu and Samet, 2002; Donaldson *et al*, 2003; Davis *et al*, 2003). The time needed for people to adopt effective responses to hotter weather is not known.

There are particular grounds for concern about mortality from occasional heatwaves of unprecedented severity (see chapter 1). In 2003, an exceptional heatwave in central France caused mean daily temperatures around 28°C for ten days with 14,000 deaths (Pirard *et al*, 2005). Further south, where people were more accustomed to high temperatures, the heatwave caused less mortality. Estimates of the probability of a heatwave approaching that level in Britain, and of its likely effect on mortality, are particularly important.

This chapter estimates heatwave related mortality using data from 1971 and from extrapolation of this until 2015:

- The level of temperature and frequency of heatwaves, and of heat related mortality in the absence of intervention.
- The changes in vulnerable population aged 65+, and the response of this population to heat and cold.

Assessments were made separately for South-East England, the rest of England and Wales, and Scotland; data for Northern Ireland were not available. An indication of longer term problems was based on current predictions of greenhouse gas emissions and climate warming over the next century is described in Chapter 1.

6.2 Observed changes in heat and cold related mortality (1971-2003)

Methods and Data

The Office of National Statistics and the Scottish Office provided populations and daily deaths 1971-2003 (1974-2003 in Scotland). Populations aged 65+ for years 1971, 1976, 2003 respectively in South-East England (Greater London, Hertfordshire, Essex, Kent, East and West Sussex, Hampshire, Surrey, Berkshire, Oxfordshire, Buckinghamshire and Bedfordshire) were 2.334, 2.467, 2.693 million, in the rest of England and Wales 4.216, 4.591, and 5.589 million, and in Scotland 0.685 (for 1974), 0.697 (for 1976), 0.809 (for 2003) million. Mortalities were obtained after dividing daily deaths by estimates of daily population obtained by fitting a 3rd order polynomial to the yearly population, estimated by ONS from census data. Recent mortality data were not available for Northern Ireland in time.

Daily temperatures for Edinburgh (Royal Botanical Gardens), for South-East England (Heathrow) and for the rest of England and Wales ('Central England Temperature' – mean of Squires Gate, Lancashire; Manchester Airport; Malvern, Worcestershire; Rothamsted, Hertfordshire) were provided by the Meteorological Office. All temperatures used were 24-hour mean values.

Heat related mortality starts in the UK when mean daily temperature exceeds about 18°C (Donaldson, Keatinge and Nayha, 2002). Excess mortalities due to heat and cold each year per 10⁶ people aged 65+ were estimated respectively as the increase in mortality above and below the 3°C band of temperature at which mortality was lowest in that region that year. No adjustment was made for air pollution (Keatinge and Donaldson, 2006) or for influenza. Cold related mortality is much larger than heat related mortality, both in the UK and in the rest of Europe (Keatinge *et al*, 2000; Eurowinter Group, 1997; Healy, 2003). The band of minimum mortality was obtained from mortalities in 3°C bands calculated at successive 0.1°C increments. Summer temperatures were means for June-August, winter temperatures means for December-February, and temperature of onset of heat related mortality was the upper end of the 3°C band of minimum mortality. Changes over time (1971 to 2003) were calculated by linear regression with confidence limits.

As described in Chapter 1, the UK has experienced some warming in the last three decades. Figure 6.1 shows that mean summer temperature in South-East England rose significantly ($p < 0.05$) from 16.6 to 18.0°C between 1971 and 2003. It rose non-significantly in the rest of England and Wales from 15.2 to 16.2°C and marginally in Scotland from 14.3 °C in 1974 to 14.7 °C in 2003.

Despite the increasing temperature, the trend in annual heat-related mortality per million aged 65+ fell significantly in Scotland and non-significantly in other regions:

- South-East England from 258 in 1971 to 193 in 2003,
- rest of England and Wales from 188 to 93,
- Scotland from 125 (in 1974) to only 8 in 2003 ($p < 0.05$).


Winter temperatures tended to rise and the trend in cold-related mortality per million aged 65+ fell in all regions ($p < 0.001$):

- South-East England from 9174 to 5903,
- rest of England and Wales from 9222 to 6088,
- Scotland from 9751 in 1974 to 6166 in 2003.

The bottom row of graphs in Figure 6.1 shows that the temperature of the upper limit of the band of minimum mortality, above which mortality starts to rise with temperature, tended to increase in all regions. This, as well as the decline in heat related mortality despite warmer summers, provides evidence of improving tolerance of the population to hot weather, especially in Scotland. The figure also shows that mean summer temperatures in South-East England in 1976 and 2003 were higher than those of neighbouring years, and produced higher heat related mortalities than these neighbouring years.

Heat related deaths in 1976 (actual, not trend and not as mortality per 10⁶) totalled 4651 in all regions of Britain; 2101 were in South-East England, 2392 in the rest of England and Wales, and 158 in Scotland. Linear regression of annual heat-related deaths in South-East England showed that these tended to fall (by 69 to 548) although summer temperature rose between 1971 and 2003 and the vulnerable population aged ≥ 65 increased (from 2.334 to 2.693 million).

Figure 6.1. Mean summer and winter temperatures, heat and cold related mortalities, and upper limit of 3°C band of minimum mortality each year. Means and 95% confidence limits of the mean.



Key


Significance of slope:

* = P<0.05

** = P<0.01

*** = P<0.001

Figure 6.2. Hottest day and hottest runs of five and nine days each year. Means and 95% confidence limits for individual years.



Key

Significance of slope:

* = $P < 0.05$

** = $P < 0.01$

6.3 Observed and future changes in heatwaves and in heatwave mortality

Figure 6.2 shows that the temperature of the hottest day each year, and of the hottest five and nine consecutive days each year in South-East England and in the rest of England and Wales, all tended to increase at approximately double the rate of increase of mean summer temperature, as variability in the daily temperatures increased (see also Schar *et al*, 2004). Changes in Scotland were too small to assess.

Figure 6.3 shows daily data for the 1976 and 2003 heatwaves. Mortality rose sharply when mean daily Central England temperature exceeded 25°C for a few days, as it did in South-East England in 1976 with one day at 25.7°C followed by three at 27.45-27.7°C (mean 27.1°C over the successive four days above 25°C). During this time daily mortality continued to rise, with approximately equal increments, and fell on the day temperature fell. The heatwave of 2003 was both briefer and less intensive, producing a clear rise in mortality mostly in South-East England.

Figure 6.4 shows that deaths per day (not adjusted for the progressive rise in population aged 65+) rose sharply in the South-East when daily mean temperature (on the same day) (Heathrow) exceeded 25 °C. Even with the increase in the population aged ≥65, heat related deaths were a little lower in the years after than before 1986, particularly at the higher temperatures.

The 95% confidence limits in Figure 6.2 show the probability of a nine day heatwave averaging 27°C in South-East England as one in forty each year by year 2012, or of one in four (25%) at some time during the ten years from mid 2007. The potential mortality from such a heatwave can be estimated as follows:

There were 854 heat related deaths in South-East England during the four day heatwave in 1976, estimated as excess deaths above the 328 mean daily deaths in the 3°C band of minimum mortality that year. During those four days (when daily temperature averaged just over 27°C) daily deaths increased each day by a mean of 80.2 from a baseline of 341 mean daily deaths on the previous 14 days. If deaths had continued to increase at that rate, nine days averaging 27°C in South-East England would have caused 3726 heat related deaths there, that is, 2382 more than actual heat related deaths in the nine day heatwave starting 25th June 1976.

For 2012, this estimate of heat related deaths during a nine day heatwave in South-East England can be reduced a little, since Figure 6.4 shows that daily deaths at such temperatures were tending to decline, by around 2.1 each year, as tolerance to heat improved. This reduces the estimate to 3046, or 1702 more than heat related deaths there on the nine days starting 25th June 1976. Taking heat related deaths outside those days in the Southeast, and all year in the rest of Britain, as unchanged from 1976, with increasing tolerance to heat offsetting rising temperatures outside the nine day heatwave in the South-East, total heat related deaths in Britain throughout 2012 would be 6353.

Figure 6.3. Temperature and mortality each day in two major heatwaves in South-East England.





Figure 6.4. Crude relationship between daily deaths and mean daily temperature in South-East England. Temperatures are means of values on days 0, -1, -2 relative to mortality.

Key

+ — 1971-1986 o — 1987-2003



6.4 Discussion

Increasing tolerance to heat has so far prevented rising temperature from increasing mean heat related mortality, even in South-East England where summer temperature rose 1.4°C since 1971. Together with falls by more than a third in cold related mortality in all regions as winters grew warmer, the overall trend in combined heat and cold related mortality during climatic warming from 1971 to 2002 was beneficial.

The increasing tolerance to heat probably resulted as much from lifestyle changes, such as greater readiness to wear informal clothing and less need for physical exertion, as from physiological adaptation to heat stress, which is relatively short term. However, we estimate that the increasingly variable as well as higher summer temperatures will create, by 2012, a one in forty risk every year (a one in four risk in the decade centred on 2012) of a nine day heatwave at 27°C in South-East England. Without preventive action, this could cause more than 3000 immediate deaths with more than 6350 heat related deaths throughout that summer.

6.5 Actions to reduce heatwave mortality as summers become hotter.

The most obvious need at this level of risk is for elderly and ill people and their carers to be aware that these people are particularly vulnerable to heat stress (Flynn *et al*, 2005), and to adopt protective measures that are common in Southern Europe in hot weather. The Heatwave Plan for England already incorporates several of the following points. We recommend that, at the start of summer, the first forecast of mean daily temperature approaching 17°C in any region should trigger initial printed and broadcast warnings, and that the broadcast warnings be repeated with any forecast of mean daily temperature above 25°C.

Important components of advice are to have an electric fan available and check that windows can be opened. Shading of windows from direct sunshine, for example by outside shutters, will prevent greenhouse heating. If shading is impractical, the use of thick curtains can reduce heating of the indoor environment. Windows should be opened in the early morning, and shut if outdoor temperature rises above indoor temperature. If indoor temperature rises enough to cause heat discomfort, a fan will need to be started. Water can be sprinkled on the face, arms and clothing if necessary to reinforce evaporative cooling by sweat, particularly in people taking medical drugs that suppress sweating. If indoor humidity rises enough to prevent evaporation and cause heat discomfort, windows will need to be opened again to provide air unsaturated with water vapour. In emergencies, a cool bath or shower will provide rapid cooling. It is important for people to continue to eat meals, preferably of carbohydrate and with at least normal amounts of salt, and to drink plenty of water to maintain osmotic balance. Loss of salt and water in sweat can otherwise cause dehydration with haemoconcentration and increased risk of coronary and cerebral thrombosis (Keatinge *et al*, 1986), as well as accelerating body heating by impairing vasodilatation and sweating.

It is important that the warnings stress the importance of immediate cooling at home, rather than waiting for admission to hospital. Overall hospital admissions in England rose by only 1% in the 2003 heatwaves, but among people in London aged >75 they rose 16% (Johnson *et al*, 2005). Many of the deaths in the more severe 2003 French heatwave occurred in hospitals (Pirard *et al*, 2005).

Unlike air conditioning, these measures do not involve high energy consumption. Air conditioning does, and so can accelerate global warming. It can be kept to a minimum by effective use of these simple measures, although it will sometimes be needed where these cannot be implemented adequately, as in hospitals or old peoples' homes when only small numbers of carers are available to attend to large numbers of incapacitated people.

Scenarios of climate change beyond 2012 take into account some of the uncertainty about future levels of greenhouse gas emissions (Chapter 1), and suggest increases in the frequency of heatwaves over the next hundred years in the South-East of the UK, and less in other regions. Such temperatures have led to extensive installation of air conditioning in the United States. Despite the adverse effect of air conditioning on carbon dioxide emissions, policy on its use in Britain, together with use of other protective measures against heat stress, needs to be reviewed regularly as further temperature and mortality data become available.

References

- Basu, R. and Samet, J.M. (2002) Relation between Elevated Ambient Temperature and Mortality: A Review of the Epidemiologic Evidence. *Epidemiologic Reviews* **24**:190-202.
- Davis, R.E., Knappenberger, P.C., Michaels, P.J. and Novicoff, W.M. (2003) Changing Heat-Related Mortality in the United States. *Environ.Health Perspect.* **111**:1712-1718
- Donaldson, G.C., Keatinge, W.R. and Nayha, S. (2003) Changes in summer temperature and heat related mortality since 1971 in North Carolina, South Finland and Southeast England. *Environmental Research* **91**:91:1-7.
- Eurowinter Group: Keatinge, W.R., Donaldson, G.C., Bucher, K., Cordioli, E., Dardanoni, L., Jendritzky, G., Katsouyanni, K., Kunst, A.E., Mackenbach, J.P., Martinelli, M., McDonald, C., Näyhä, S. and Vuori, I. (1997) Cold exposure and winter mortality from ischaemic heart disease, cerebrovascular disease, respiratory disease, and all causes in warm and cold regions of Europe. *The Lancet* **349**:1341-1346.
- Greenberg, J.H., Bromberg, J., Reed, C.M., Gustafson, T.L. and Beauchamp, R.A. (1983) The epidemiology of heat-related deaths, Texas - 1950, 1970-79, and 1980. *Am.J.Public Health* **73**:805-7.
- Healy, J.D. (2003) Excess winter mortality in Europe: a cross country analysis identifying key risk factors. *J.Epidemiol.Comm.Health* **57**:784-789.
- Johnson, H., Kovats, R.S., McGregor, G., Stedman, J., Gibbs, M., Walton, H., Cook, L. and Black, E. (2005) The impact of the 2003 heat wave on mortality and hospital admissions in England. *Health Statistics Quarterly* **25**:6-11.
- Keatinge, W.R., Coleshaw, S.R.K., Easton, J.C., Cotter, F., Mattock, M.B. and Chelliah, R. (1986) Increased platelet and red cell counts, blood viscosity and plasma cholesterol level during heat stress, and mortality from coronary and cerebral thromboses. *American Journal of Medicine* **81**:795-800.
- Keatinge, W.R., Donaldson, G.C., Cordioli, E., Martinelli, M., Kunst, A.E., Mackenbach, J.P., Nayha, S. and Vuori, I. (2000) Heat related mortality in warm and cold regions of Europe, observational study. *Brit.Med.J.* **321**:670-3.
- Keatinge, W.R. and Donaldson, G.C. (2006) Heat acclimatization and sunshine cause false indications of mortality due to ozone. *Environmental Research* (in press).
- Pirard, P., Vandentorren, S., Pascal, M., Laaidi, K., Le Tertre, A., Cassadou, S. and Ledrans, M. (2005) Summary of the mortality impact assessment of the 2003 heat wave in France. *Euro Surveill.* Jul 1;10 (Epub ahead of print).
- Rogot, E., Sorlie, P.D. and Backlund, E. (1992) Air-conditioning and mortality in hot weather. *American Journal of Epidemiology* **136**:106-116.
- Taylor, A.J. and McGwin, G. (2000) Temperature-related deaths in Alabama. *South Med.J.* **93**:787-92.

7 The health impact of climate change due to changes in air pollution

(See also section 4.7 of 2001 report)

HR Anderson, St George's, University of London
RG Derwent, Rd Scientific
JR Stedman, AEA Technology
GD Hayman, AEA Technology

Summary

- Future ambient concentrations of air pollutants which are hazardous to health will depend both on trends in emissions of primary and precursor pollutants and trends in meteorological factors which affect dispersal of pollutants and secondary chemical atmospheric processes.
- The weather conditions associated with winter episodes will become less common while over the same time there will be a decrease in the ambient concentrations of PM₁₀, SO₂ and NO₂ should an episode occur. O₃ concentrations are already low in winter episodes and will remain so in any future episodes. Thus the health impact of winter episodes, which is already low, will become even less.
- Weather conditions associated with summer episodes will increase in frequency and intensity. However, because the European emissions of ozone precursors such as VOCs and NO_x are predicted to fall, episodic peak ozone concentrations should continue to fall over the period to 2030. The frequency and intensity of summer NO₂ episodes is unlikely to change. PM₁₀ usually increases during summer episodes, but the intensity of this is likely to fall in the future because of falling emissions. SO₂ episodes, already rare, will not increase.
- Longer term mean concentrations of particles, NO₂ and SO₂ are unlikely to increase in the future. However, annual mean and annual mean daily maximum 8-hour ozone concentrations are likely to increase in both the short term until 2030 and over the long term until 2100 as part of a more general increase in ozone occurring in the hemisphere as a result of both increases in precursor emissions and climate change.
- Without the assumption of a threshold, the increase in estimated annual ozone concentrations between 2003 and 2020 will result in a 15% increase in attributable deaths and hospital admissions for respiratory diseases. When threshold assumptions of 35 ppb and 50 ppb (8-hour average) are assumed, the respective increases are 51% and 53%. Estimates holding climate at 2003 show increases of 8%, 14% and -5% for the no threshold, and 35 ppb and 50 ppb thresholds respectively. Thus, in the no threshold model, about half of the increase in health impact is due to changes in UK and European emissions rather than climate change. In the threshold models however, most of the increase is attributable to climate change.

Recommendations

- Maintain and strengthen measures that reduce outdoor air pollution.

7.1 Introduction

Urban air quality in the United Kingdom has been extensively reviewed in recent years, together with its public health consequences. Understanding has also grown concerning the important linkages between urban air quality and global climate change. There is growing concern that some of the steps taken to tackle urban air quality may either exacerbate or ameliorate climate change. Equally well, climate change and the global scale population and economic growth that drives it, may make urban air quality goals more difficult to achieve in the medium term (Air Quality Expert Group, 2006).

Here we look at the period up to 2030 and beyond and consider how the emissions of the classic air pollutants may change in the United Kingdom. We use a global climate model to understand how climate change may influence the frequency and intensity of the meteorological conditions associated with pollution episodes. By combining these two elements, we can estimate future levels of ambient air pollutant concentrations. Public health impacts have been assessed using risk coefficients based on recent epidemiological evidence. The air pollutants of particular interest are particulate matter (PM₁₀), sulphur dioxide (SO₂), nitrogen dioxide (NO₂), and ozone (O₃).

In the previous report on climate change and public health in the United Kingdom, we presented estimates of the human health impacts of air pollution through to the year 2100 (Anderson, Derwent, & Stedman, 2000). In the intervening period, several aspects of our methodology and approach have been subject to review and updating. Furthermore, the policy focus has shifted more towards the period up to the year 2030. A reassessment of the human health impacts of air pollution and climate change in the United Kingdom is therefore timely.

7.2 Meteorological factors, emissions, and future urban air pollution episodes

Meteorological factors

As explained in our previous report, to address the occurrence of the meteorological conditions which drive wintertime air pollution episodes, we have examined the daily meteorological parameters produced by the Hadley Centre Climate Model (HadCM3) for each day through to December 2099. This is a coupled ocean-atmosphere global climate model, driven by the increasing concentrations of greenhouse gases and aerosols as described by the IPCC IS92a (Business as usual) scenario (IPCC 1996). The output has been taken for a single grid point in the British Isles, [52.5oN, 0oW], which corresponds with the location of Central England.

We have looked at the simultaneous occurrence of low windspeed with freezing conditions that is associated with episodes of poor urban air quality. There are isolated days with low windspeeds and below freezing temperatures during the decades up to the 2030s, but their frequency is reduced from the 2050s onwards. The number of days with temperatures just below zero and low windspeeds also declines dramatically through to the year 2100. The conditions of low windspeeds and below freezing temperatures that have occurred with a frequency of about once per decade during the second half of the 20th century, appear much less frequently as winters gradually become warmer and windier. On this basis we concluded that wintertime pollution episodes were likely to become less frequent and less intense in the future.

From time to time during warm, sunny anticyclonic conditions, elevated ozone concentrations occur over the British Isles as hazy, regionally-polluted air masses drift in from continental Europe. Elevated ozone concentrations tend only to be recorded

when daily temperatures exceed 25°C and when winds are light and from the East. To address the occurrence of the meteorological conditions which drive these summertime air pollution episodes, we have again examined the daily meteorological parameters produced by the Hadley Centre Climate Model (HadCM3). It is the simultaneous occurrence of daily maximum temperatures above 25°C and low windspeed conditions which favour the occurrence of summertime air pollution episodes and so the joint distribution of elevated temperatures and low windspeed conditions has been investigated.

There are an increasing number of days with low windspeeds and daily temperatures above the 25°C threshold for elevated ozone levels for each of the decades through to the year 2100. They might triple in frequency by the 2030s and increase by an order of magnitude by the 2080s. An increasing number of these low windspeed days have maximum temperatures above 35°C by the 2050s and above 40°C by the 2070s. Conditions of low windspeeds and elevated daily temperatures appear to occur with increasing frequency throughout the period up to 2030 and beyond, driving up the frequency of conditions associated with summertime air pollution episodes. On this basis, we asserted that ozone episodes had the potential to become more frequent and more intense in the future.


Since the previous study, high resolution data has become available from a regional climate model HadRM3P and this has been used as a basis for the UKCIP02 climate scenarios discussed elsewhere in this report (UKCIP02, 2002). These data are available at a spatial resolution of 50 km x 50 km and this represents a significant improvement on the single grid point data provided by the global model HadCM3. However, we have not updated our assessments of the frequency and intensity of wintertime and summertime air pollution episodes from those based on the global model to those based on the regional model. This is because of the poor performance of the HadRM3P model during the control period from 1960 to 1991 for the parameters required in the evaluation of urban air quality. Due to the excessive drying out of soil surfaces, almost one in every two summers in the Southeast of England were found to have maximum temperatures in excess of 35°C with a maximum temperature of over 41°C (Moberg & Jones, 2004). Such behaviour would not provide a satisfactory base case for a regional air quality model.

Future urban air pollution episodes

Emissions of primary pollutants in urban areas from road traffic are expected to decline steadily from current levels over the period to 2030 and beyond due to the continuing tightening of both fuel and vehicle emission legislation. Figures 7.1 – 7.3 show the anticipated falls in traffic emissions of PM₁₀, oxides of nitrogen (NO_x) and volatile organic compounds (VOCs), respectively from their peak levels in the 1990s by 2025. If this were the only factor operating, then urban PM₁₀ air quality would steadily improve through to 2025. Taken together with the tendency for winters to become warmer and windier, then both the frequency and intensity of wintertime PM₁₀ pollution episodes should decrease during the period up to 2025.

Wintertime episodes of elevated SO₂ and black smoke concentrations have in the past been confined to the areas where coal has been the most important domestic fuel. But because this use of coal has declined markedly in British cities, such episodes are of historic importance only. Summertime episodes of SO₂ are generally associated with plumes from individual major industrial plants and have not been considered in any detail here. However, the current trend towards tighter regulation, cleaner fuels and the implementation of cleaner technologies in industries such as power generation, should lead to a reduction in the frequency of summertime episodes of SO₂ by 2025.

Wintertime episodes of NO₂ depend for their occurrence upon the chemical reaction:



Because this mechanism depends on the square of the nitric oxide (NO) concentration, wintertime episodes are expected to become increasingly rare as NO_x emissions decline steadily (see Figure 7.2), from their peak in the early 1990s and as the frequency of wintertime stagnant conditions declines with climate change.

Episodes of elevated NO₂ concentrations occur also in the summertime when photochemical episodes occur simultaneously with episodes of poor dispersion of NO_x emissions. The intensity of summertime NO₂ episodes is expected to decline over the period out to 2025 as a result of both the reductions in urban NO_x emissions from road traffic, see Figure 7.2, and the expected decline in episodic ozone concentrations resulting from reductions in European regional emissions of ozone precursors. However, the frequency of summertime episodes may increase at the same time due to climate change. Furthermore, there is also the impact of the increase in the global ozone baseline to consider as well as the increasing direct emissions of NO₂ into urban atmospheres from diesel vehicles fitted with catalytically regenerating particle traps and filters. At present, these factors are finely balanced and so it is likely that there will be little change in the intensity and frequency of summertime NO₂ episodes for the present. A detailed study of future NO₂ concentrations in the United Kingdom is available elsewhere (Air Quality Expert Group, 2004).

Figure 7.1. UK urban road traffic PM₁₀ emissions up to 2030.




Figure 7.2. UK urban road traffic NO_x emissions up to 2030.





Figure 7.3. UK urban road traffic VOC emissions up to 2030.



Future ozone episodes

It is recognized that the summertime European regional scale photochemical ozone episodes sit on top of an ozone baseline which is itself global in scale. Ozone monitoring carried out at the Montsouris Observatory near Paris during the second half of the 19th century shows ozone levels which are about one half to one third of present-day ozone levels (Volz & Kley, 1988). Computer modelling studies have shown that this apparent increase in Northern hemisphere baseline ozone levels from pre-industrial times through to the present day could have been caused by increasing emissions of methane (CH₄), CO and NO_x from human activities (IPCC, 2001). Furthermore, these computer modelling studies suggest that this ozone increase should continue into the future. In our previous report, the need was recognized for further modelling at the global and regional scale of the combined effects of climate change and changes in the emissions of the precursors of both the background and regional ozone in the period up to 2030.

More recently, further global emission scenarios have been developed for the tropospheric O₃ precursor gases, in addition to those used previously and promulgated by the Intergovernmental Panel on Climate Change in their SRES scenarios. These new global emission scenarios have prompted a further consideration of the combined influence of ozone precursor emission controls on both global and regional ozone within Europe. They address the global emissions of man-made sources of SO₂, NO_x, CH₄, CO and VOCs annually through to 2030 in two scenarios: CLE, current legislation and MFR, maximum feasible reduction. In the CLE scenario, each country applies the pollution controls that have already been agreed with their current policies, whereas in the MFR scenario, all technically and economically feasible ozone precursor emission reductions are put in place by 2030.


Figure 7.4 presents the annual mean ozone concentrations for the Southeast of England calculated with the global three-dimensional Lagrangian chemistry-transport model STOCHEM from 1990 through to 2030 (Derwent *et al*, 2006). Model results are shown as solid black lines and the available observations are taken from the DEFRA Rural Ozone Network. The observations are presented as the annual average of the daily maximum 8-hour mean ozone concentrations. Over the period from 1990 through to 2003, the model and observations agree well both in relative magnitudes and trends. The observations show a small downwards trend of -0.03 ppb/year and the model results between +0.27 and -0.10 ppb/year depending on the climate dataset employed. In the global scenario with all countries, both within Europe and globally, implementing their current legislation CLE, ozone levels increase steadily over the period to 2030 by +0.10 to +0.13 ppb/year. When all countries adopt maximum feasible reductions, ozone levels decline in Southeast England by -0.04 ppb /year. Relative to 2000 levels, ozone concentrations rise by about 3.0 ppb by 2030 in the CLE scenario and decline by about 1.4 ppb in the MFR scenario.

The ozone concentration changes anticipated in the CLE and MFR scenarios in Figure 7.4 by the year 2030, are somewhat smaller than the 3.7 ppb increase reported in our previous assessment using the IPCC SRES A2 scenario. This reflects a change in perceptions of the growth in Asian emissions and the take-up of ozone precursor emission controls in the same region that have been built in to the new emission scenarios. However, on the longer timescale, the ozone increases projected in our previous report with the IPCC SRES A2 scenario by 2060 and 2100 still remain a distinct possibility. Furthermore, there is no guarantee that the countries of the world will embrace the concept of maximum feasible reductions and fully implement them and so bring down both global and regional ozone concentrations as shown in Figure 7.4. Currently, therefore, the CLE curves in Figure 7.4 represent our best estimate of where future ozone concentrations will be over the period to 2030.

In the IPCC SRES A2 scenario, ozone concentrations are projected to rise throughout the 21st century. There is an important question as to the extent to which these ozone

increases are the direct consequences of climate change or the consequences of the increasing global emissions of the radiatively-active trace gases that drive climate change. Because of the close coupling between the climate processes, atmospheric chemistry and the biogeographical cycles of the radiatively-active trace gases, it is difficult to give a simple answer to these questions. Broadly speaking, ozone concentrations over the time horizon out to 2030 and 2050 are more strongly influenced by the increasing emissions of CH₄, NO_x, CO and VOCs that drive climate change. In contrast, over the time horizon from 2030 onwards, climate change processes themselves start to exert a more dominant influence on future ozone concentrations. Nevertheless, in a given snap-shot in time or over a given timescale, it is difficult to discern whether climate change itself or increasing global emissions of methane are responsible for the ozone changes from the present day because the global systems are so closely coupled.

Figure 7.4 *Model calculated monthly mean ozone concentrations for Southern England from 1990 through to 2028 in two global emissions scenarios: triangles, on current emission control policies (CLE) and crosses, with maximum feasible emission reductions in methane, CO, NO_x and VOCs (MFR). The blue trace shows the observed annual means of the daily 8-hour mean ozone concentrations from the rural site network also in Southern England.*



7.3 Health impacts: episodes

We have not attempted to quantify the changes in health impact associated with the trends in the frequency and atmospheric characteristics of winter and summer episodes described above. One reason is that there is uncertainty about what concentrations might occur. Another reason is that there is little firm evidence concerning the appropriate exposure response coefficients to apply during episode conditions. Finally, since episodes account for such a small proportion of days, their impact is far outweighed by the effects of day-to-day exposure to air pollution during the rest of the year. Nevertheless, air pollution episodes are important public health threats and as such are associated with publicity and the issuing of advice to the public. Qualitatively the following predictions can be made.

The weather conditions associated with winter episodes will become less common while over the same time there will be a decrease in the ambient concentrations of PM₁₀, SO₂ and NO₂ should an episode occur. O₃ concentrations are already low in winter episodes and will remain so in any future episodes. Thus the health impact of winter episodes, which is already low, will become even less due to the combined effects of climate change and reduced emissions.

In contrast, the weather conditions associated with intense summer-time photochemical ozone episodes will increase in frequency. However, these future episodes will be accompanied by lower concentrations of ozone than already occur under similar circumstances because European emissions of VOCs and NO_x precursors are predicted to fall.

We have concluded that the frequency and intensity of summer NO₂ episodes is unlikely to change. The implications of this for health impact are uncertain partly because there is more doubt about the direct health effects of NO₂ than for particles and ozone (WHO, 2004). PM₁₀ usually increases during summer episodes, but the intensity of this is likely to fall in the future. SO₂ episodes, already rare, will not increase.

7.4 Quantitative health impact assessment

The main pollutant of concern is O₃, concentrations of which are predicted to rise in the long-term, largely due to increases in hemispheric ozone. In this section we estimate changes in attributable deaths and emergency respiratory hospital admissions attributable to this trend up to 2020. This was done in the previous report but needs repeating because of changes in the various parameters required to estimate health impact (predicted exposure to O₃, risk coefficients, baseline rates, and population estimates). The choice of exposure responses coefficients is not clear cut. Since the first climate change report, there have been important refinements in epidemiological methods (Health Effects Institute, 2003) and an expansion in the body of evidence associating ozone with health effects, including several multi-city studies (Department of Health, 2006, in preparation).

Choice of mortality coefficient for ozone

Unlike the case of particles, for ozone there is insufficient evidence for an effect of chronic exposure on mortality to justify estimating health impacts in terms of shortened life-expectancy. However, mortality associations have been found in a large number of short term exposure studies using daily time-series. These enable the estimation of attributable deaths “brought forward”, but the life lost cannot be quantified. A variety of averaging times (1 hr, 8-hr and 24-hr) have been used in these studies but we shall

confine our estimates to the 8-hour maximum ozone. This has been used in many European time-series studies, and was recently adopted by the WHO in its forthcoming update of guidelines. The previous climate change report used a coefficient of 0.6% increase in daily deaths per $10\mu\text{g}/\text{m}^3$ 8-hour ozone. This was taken from the QUARK 1 report (Department of Health Committee on the Medical Effects of Air Pollutants, 1998) and was based on unpublished results from the European multi-city study (APHEA 1) which had combined results from 5 cities (Athens, Barcelona, London, Lyon and Paris) (Touloumi *et al*, 1997). Many new studies have been reported as well as several meta-analyses.

A meta-analysis of 15 European cities carried out by WHO (2004), has been used to estimate mortality impacts of ozone by Clean Air For Europe (CAFE) for the European Commission (Commission of the European Communities, 2005). This estimate was 0.3% (95% confidence interval 0.1 to 0.4). It did not incorporate the more recently published results from 23 European cities as reported from APHEA 2. (Gryparis *et al*, 2004) While there was overlap between the cities included in the APHEA 2 and WHO meta-analyses, the APHEA 2 analysis had the advantage of newer data sets and a standardised and more refined approach to the analysis. To obtain the most recent and comprehensive meta-analysis of European cities, we combined these new European results with those from four additional European cities (27 cities in total) and obtained a random effects estimate of 0.3% (95% CI -0.05 to 0.74), the same as that obtained by the WHO meta-analysis. This is also what was obtained by the COMEAP review of ozone (Department of Health, 2006 - in preparation) in a meta-analysis based on 22 cities world-wide (0.3%, 95% CI 0.2 to 0.4). Since three independent meta-analyses with different mixes of cities all agree, we decided to use the all-year coefficient for 8-hour ozone of (0.3% 95% CI 0.1 to 0.4) based on European cities published by WHO and used by CAFE in its health impact assessment for the European Commission (WHO, 2004). It is within the range of estimates provided by three recent meta-analyses reported from the US based on 1 hour and 24-hour averaging times (Bell, Dominici & Samet, 2005; Ito, De Leon & Lippmann, 2005; Levy, Chemerynski & Sarnat, 2005). This coefficient is about half of that used in the previous climate change report.

Choice of morbidity coefficient for ozone

Ozone has been associated with a range of morbidity outcomes, including emergency hospital admissions for respiratory disease, increased respiratory symptoms in asthmatics and reduced lung function (WHO, 2003). Notably, however, there is little evidence of an association with hospital admissions for heart disease or stroke admissions (Department of Health Committee on the Medical Effects of Air Pollutants, 2006). Hospital admission is the only morbidity outcome for which there is adequate baseline data and we have therefore confined our morbidity impact assessment of ozone to emergency respiratory admissions.

The previous report estimated the impact of ozone on emergency respiratory admissions using a coefficient of 0.7%, based on a rounded estimate from unpublished results from four European cities in the APHEA 1 study (Spix *et al*, 1998). No results for 8-hr ozone and all-age all respiratory admissions are yet available from APHEA 2. However, newer estimates are available from individual city studies. We combined the results of all available European cities (Birmingham, London, Paris and Rome) and obtained an estimate of 0.1% (95% CI -0.3 to 0.6). The WHO meta-analysis of European cities obtained age specific estimates for the 15-64 age-group (1.001, 0.991 to 1.012) and 65+ age-group (1.005, 0.998 to 1.012). These age-specific coefficients provide an alternative for estimating attributable admissions which has the advantage of being derived from the same WHO meta-analysis as the mortality coefficient. We shall estimate impacts using both the coefficient for all ages and the WHO age-specific coefficients.

Most studies show higher coefficients in the warm than the cool half of the year. For this reason, health impacts are frequently estimated for the warm half of the year only (Stedman *et al*, 1997). This approach was also followed by QUARK 1, and the previous

Climate Change report, using the all year coefficients, which is likely to have underestimated the exposure response. In contrast, the APHEIS/ENHIS project used larger warm-season-specific coefficients for estimating health impacts of ozone (ENHIS, 2005). The argument against using all-year coefficients is based mainly on the belief that there is likely to be a threshold for ozone effects together with the possibility that negative confounding with traffic pollution in the cool season may artificially depress ozone coefficients based on cool season analyses. Another argument is that individual ozone exposure is greater in the summer because more time is spent outdoors and an increased outdoor to indoor ratio. Against this argument is the general lack of epidemiological evidence of a threshold where this has been looked for in time-series analyses. Having considered these arguments we decided to estimate the health impacts using all-year ozone estimates as was also recommended by UNECE (UNECE, 2004).

Because the question of a threshold for ozone remains open, it is usual to examine the sensitivity of the results to a threshold assumption. This was done in a study of the effect of ozone in the summer of 1995 (Stedman, Anderson, Atkinson & Maynard, 1997), in the previous climate change report (Anderson, Derwent & Stedman, 2000) and in Quark 1 (Department of Health Committee on the Medical Effects of Air Pollutants, 1998). In these analyses, a threshold of 50 ppb 8-hr ozone was used to estimate health effects on days at or above that level. More recently, the question of a threshold for ozone was discussed by the Joint Task Force on the Health Aspects of Air Pollution (UNECE, 2004). While it was recognised that current evidence was insufficient to derive a threshold, it was also thought that uncertainty about exposure response increased at lower levels. This led to the decision to estimate effects only in relation to days over 35 ppb, this being about the background level in Europe. This cut off was adopted by the CAFE impact assessment. The same group also recommended that effects should be estimated over the whole year using all -year coefficients. Recognising that this approach might underestimate effects it was also recommended that a sensitivity analysis using no threshold should be also done. Since urban monitors had been used to generate exposure response relationships, urban levels of ozone should be used to estimate urban health effects. We estimated impacts with no threshold, a threshold of 35 ppb and a threshold of 50 ppb (as was done in the previous climate change report).

Ozone modelling

The Ozone Source Receptor Model (OSRM) has also been used to generate UK scale maps on a 10 km x 10 km grid. The OSRM is a recently developed model to describe photochemical ozone production in the UK (Hayman *et al*, 2004; Hayman *et al*, 2005; Hayman, Jenkin & Pilling, 2002). The OSRM covers the EMEP model domain and uses global meteorological datasets provided by the Met Office to derive 96-hour back trajectories. The chemical scheme is based on that used in the STOCHEM model (Collins *et al*, 1997). The mechanism has approximately 70 chemical species involved in approximately 180 thermal and photochemical reactions. The mechanism represents ozone formation using 10 VOCs, which provides an appropriate description of ozone formation on the regional scale. The emission inventories are taken from EMEP for Europe with the option to use NAEI emission inventories for the UK, which have been aggregated to 10 km x 10 km and into 8 key sectors.

The OSRM describes the boundary layer by a single box and assumes that this is well mixed. While this is a reasonable assumption for rural areas, it is less valid in urban areas with high NO_x emissions. In such areas, there will inevitably be a gradient in the NO_x concentration profile, with higher concentrations at the surface. This will lead to lower ozone concentrations at the surface through reaction with NO.

An algorithm has been developed and implemented in the OSRM post-processor to convert the mid-boundary layer concentrations to surface concentrations on a hour-by-hour basis. This algorithm uses the local meteorological parameters characterising the boundary layer, surface roughness appropriate for the surface types considered,

resistance parameters for O₃ and NO₂, the local NO_x emission rates and a simple NO-NO₂-O₃ photostationary state chemistry.

The OSRM calculates ozone concentration on a hourly basis and can thus be used to calculate a wide range of ozone metrics including the annual means of the daily maximum running 8-hour mean concentrations with cut-offs at 0, 35 and 50 ppb used here. A comparison of the OSRM results for these metrics with monitoring data and empirical modelling methods has been provided by Kent *et al* (Kent *et al*, 2005).

Estimation of attributable deaths and admissions

In calculating the effects of pollutants on health the following sequence of steps has been adopted. These steps are described in more detail in earlier reports (Department of Health Committee on the Medical Effects of Air Pollutants 1998; Stedman *et al*, 2002).

- a. The country has been divided into 1 km grid squares and the concentration of pollutants and resident population has been estimated for each. The former has been derived from the 10 km square OSRM results and the latter from 2001 census data. Population-weighted mean concentrations are then calculated by region or for the whole of the UK.
- b. A baseline level of the given health-related and pollution affected events e.g. daily deaths, hospital admissions for the treatment of respiratory diseases was obtained from National Statistics and the Hospital Episode System (England only).
- c. By combining the data from (a) and (b) and applying the chosen coefficient linking the pollutant concentration with the relevant effects, the expected health effects were derived under various threshold assumptions (none, 35 ppb and 50 ppb).
- d. To distinguish effects of climate and hemispheric emissions from those of trends in UK emissions, estimates were made for 2020 under the climate change scenario, and under the UK and European emissions scenario holding the climate and hemispheric emissions at 2003.

Results

Table 7.1 shows the results of modelling the health impacts of predicted changes in ozone using the above approach. Table 7.2 summarizes these results and also presents the qualitative conclusions concerning other pollutants and winter and summer episodes.

The percentage change in ozone concentration and thus the health impact between 2003 and 2020 is dependent on both the ozone metric chosen and the assumptions about the composition of the atmosphere. The ozone metric with no threshold shows a 15% increase to 2020 for the baseline and an 8% increase to 2020 for the 2003 atmosphere calculation. Thus, about half of the expected change is due to the changing atmosphere (hemispheric emissions and climate change) and half is due to changes in UK emissions, specifically the control of NO_x emissions, which is expected to lead to an increase in ozone concentration in urban areas to more closely resemble those in rural areas. There is a much larger expected increase of about 50% in the 35 ppb and 50 ppb ozone metrics to 2020 for the baseline. Most of this is due to the changing atmosphere. Reductions in emissions lead to a decrease in the 50 ppb ozone metric for the 2020 baseline with the 2003 atmosphere. For this metric the reduction in the emission of the precursors of photochemical episodes between 2003 and 2020 outweighs the increase in ozone concentration in urban areas (Hayman, Thompson, Abbott & Bush, 2005).

Table 7.1. 2003 baseline mortality and emergency hospital admissions and predictions for 2020 under (1) model for climate change including UK emissions, (2) model for UK and European emissions without climate change.

	2003 (base year)	2020 baseline (climate change, and UK emissions)	2020 baseline (2003 climate, 2020 UK and European emissions)
No threshold			
Deaths (0.3%) (95% CI 0.1 to 0.4))	11272 (3757 to 15029)	12930 (4310 to 17240)	12140 (4047 to 16186)
Resp Ads (0.1% (95% CI -0.3 to 0.6)	3664 (-10991 to 21983)	4203 (-12608 to 25216)	3946 (-11838 to 23675)
Resp Ads (sum of age specific)	9280 (-12875 to 32501)	10645 (-14769 to 37282)	9995 (-13867 to 35004)
35ppb threshold			
Deaths (0.3%) (95% CI 0.1 to 0.4))	1582 (527 to 2109)	2391 (797 to 3189)	1802 (601 to 2402)
Resp Ads (0.1% (95% CI -0.3 to 0.6	514 (-1542 to 3085)	777 (-2332 to 4664)	586 (-1757 to 3513)
Resp Ads (sum of age specific)	1302 (-1807 to 4561)	1969 (-2732 to 6896)	1483 (-2058 to 5195)
50ppb threshold			
Deaths (0.3%) (95% CI 0.1 to 0.4))	333 (111 to 444)	511 (170 to 681)	317 (106 to 423)
Resp Ads (0.1% (95% CI -0.3 to 0.6	108 (-325 to 650)	166 (-498 to 996)	103 (-309 to 619)
Resp Ads (sum of age specific)	274 (-381 to 961)	420 (-583 to 1472)	261 (-362 to 915)

Table 7.2. Summary of predictions. The range against hospital admissions refers to the two coefficients used.

	2020. Climate change model (climate, hemispheric emissions and UK emissions)	2020. UK emissions but 2003 climate	Winter episodes	Summer episodes
Pollutant	% change in deaths from 2003 (n)	% change in deaths from 2003 (n)		
Particles			Reduction	Reduction
Ozone (no threshold)			Not applicable	Increase in frequency. Lower intensity
Deaths	15% (1658)	8% (868)		
Respiratory admissions	15% (539 to 1365)	8% (282 to 715)		
Ozone (35 ppb threshold)				
Deaths	51% (810)	14% (220)		
Respiratory admissions	51% (77 to 667)	14% (263 to 667)		
Ozone (50 ppb threshold)				
Deaths	53% (177)	-5% (-16)		
Respiratory admissions	53% (58 to 146)	-5%. (-5 to -13)		
Nitrogen dioxide			Reduction	No change in frequency or intensity
Sulphur dioxide			Reduction	Reduction

Acknowledgements

Stephen Bremner and Richard Atkinson (George's, University of London) for providing meta-analyses of health risk coefficients from the Air Pollution Epidemiology Database (APED). Ramyani Gupta and Liz Limb (St George's, University of London) for supplying base rates for mortality and hospital admissions from the Lung and Asthma Information Agency (LAIA).

References

- Air Quality Expert Group. (2004) *Air quality expert group: Nitrogen dioxide in the United Kingdom* DEFRA, London.
- Air Quality Expert Group. (2006) *Air quality and climate change: a UK perspective*, DEFRA, London.
- Anderson, H.R., Derwent, R.G. & Stedman, J.R. (2000) "Air pollution and climate change," in *The health effects of climate change in the UK*, Department of Health, London, pp. 193-217.
- Bell, M.L., Dominici, F. & Samet, J.M. (2005) A meta-analysis of time-series studies of ozone and mortality with comparison to the national morbidity, mortality, and air pollution study1. *Epidemiology* **16**;4:436-445.
- Collins, W.J., Stevenson, D.S., Johnson, C.E. & Derwent, R.G. (1997) Tropospheric ozone in a global-scale three dimensional Lagrangian model and its response to NOx emission controls. *Journal of Atmospheric Chemistry* 26:223-274.
- Commission of the European Communities (2005) *Thematic Strategy: communication from the Commission to the Council and the European Parliament*.
- Department of Health Committee on the Medical Effects of Air Pollutants. (1998) *Quantification of the effects of air pollution on health in the United Kingdom*, HMSO, London.
- Department of Health Committee on the Medical Effects of Air Pollutants. (2006) *Air Pollution and Cardiovascular Disease*.
- Derwent, R.G., Simmonds, P.G., O'Doherty, S., Stevenson, D.S., Collins, W.J., Sanderson, M.G., Johnson, C.E., Dentener, F., Cofala, J., Mechler, R. & Amann, M. (2006) External influences on Europe's air quality: baseline methane, carbon monoxide and ozone from 1990 to 2030 at Mace Head, Ireland. *Atmospheric Environment* **40**:844-855.
- ENHIS. (2005) *WP5 Health Impact Assessment. Full report on HIA of outdoor air pollution*.
- Gryparis, A., Forsberg, B., Katsouyanni, K., Analitis, A., Touloumi, G., Schwartz, J., Samoli, E., Medina, S., Anderson, H.R., Niciu, E.M., Wichmann, H.E., Kriz, B., Kosnik, M., Skorkovsky, J., Vonk, J.M. & Dortbudak, Z. (2004) Acute effects of ozone on mortality from the "Air pollution and health: a European approach project1", *Am.J.Respir.Crit.Care Med.* **170**;10:1080-1087.
- Hayman, G.D., Bush, A., Kent, A., Derwent, R.G., Jenkin, M.E., Pilling, M.J. & Abbott, J. (2004) *Modelling of tropospheric ozone. The first annual report produced for the Department for Environment, Food and Rural Affairs and Devolved Administrations on Contract EPG 1/3/200*.
- Hayman, G.D., Jenkin, M.E. & Pilling, M.J. (2002) *Modelling tropospheric ozone formation. A final project report produced for the Department for Environment, Food and Rural Affairs and Devolved Administrations on Contract EPG 1/3/143*.
- Hayman, G.D., Thompson, C.L., Abbott, J. & Bush, T. (2005) *Ozone modelling for the review of the air quality strategy. Report (AEAT/ENV/R/2092 issue 1) prepared for the Department for Environment, Food and Rural Affairs and Devolved Administrations, The Scottish Executive, The National Assembly for Wales and the Department of Environment in Northern Ireland*.
- Health Effects Institute. (2003) *Revised analyses of time-series studies of air pollution and health. Special Report*. Health Effects Institute, Boston MA.

- IPCC. (1996) *IPCC second assessment report. Climate change 1996: the scientific basis* Cambridge University Press, New York.
- IPCC. (2001) *IPCC third assessment report. Climate change 1996: the scientific basis* Cambridge University Press, New York.
- Ito, K., De Leon, S.F. & Lippmann, M. (2005) Associations between ozone and daily mortality: analysis and meta-analysis1. *Epidemiology* **16**;4:446-457.
- Kent, A., Thompson, C.L., Grice, S. & Stedman, J.R. (2005) *Comparison of health based metrics and modelling methods for ozone in the UK National Atmospheric Emissions Inventory. Report AEAT/ENV/R/1935 issue 2*, AEA Technology, National Environmental Technology Centre.
- Levy, J.I., Chemerynski, S.M. & Sarnat, J.A. (2005) Ozone exposure and mortality: an empiric bayes metaregression analysis. *Epidemiology* **16**;4:458-468.
- Moberg, A. & Jones, P.D. (2004) Regional Climate model simulations of daily maximum and minimum near-surface temperatures across Europe compared with station data 1961-1990", *Climate Dynamics* **23**;695-715.
- Spix, C., Anderson, H.R., Schwartz, J., Vigotti, M.-A., Le Tertre, A., Vonk, J., Touloumi, G., Balducci, F., Piekarski, T., Bacharova, L., Tobias, A., Ponka, A. & Katsouyanni, K. (1998) Short-term effects of air pollution on hospital admissions of respiratory diseases in Europe: a quantitative summary of APHEA study results. *Arch.Environ.Health* **53**:54-64.
- Stedman, J.R., Anderson, H.R., Atkinson, R.W. & Maynard, R.L. (1997) Emergency hospital admissions for respiratory disorders attributable to summer time ozone episodes in Great Britain [see comments]. *Thorax* **52**;11:958-963.
- Stedman, J.R., King, K., Holland, M. & Walton, H. (2002) *Quantification of the health effects of air pollution in the UK for revised PM10 objective analysis. Report AEAT/ENV/R/1162*, AEA Technology Environment, National Environmental Technology Centre.
- Touloumi, G., Katsouyanni, K., Zmirou, D., Schwartz, J., Spix, C., de Leon, A. P., Tobias, A., Quennel, P., Rabaczko, D., Bacharova, L., Bisanti, L., Vonk, J. M., & Ponka, A. (1997) Short-term effects of ambient oxidant exposure on mortality: a combined analysis within the APHEA project. *Air Pollution and Health: a European Approach. Am.J.Epidemiol.* **146**;2:177-185.
- UKCIP02 (2002) *Climate change scenarios for the United Kingdom: the UK CIP02 Scientific Report*, Tyndall Centre, School of Environmental Sciences, University of East Anglia.
- UNECE (2004) *Modelling and assessment of the health impact of particulate matter and ozone. Summary report prepared by the Joint Task Force on health aspects of long-range transboundary air pollution of the WHO/ECEH. Bonn, 6-7 May 2004.*
- Volz, A. & Kley, D. (1988) Evaluation of the Montsouris series of ozone measurements made in the 19th century. *Nature* **332**;(6161):240-242.
- WHO. (2003) *Health aspects of air pollution with particulate matter, ozone and nitrogen dioxide. Report on a WHO working group. EUR/03/5042688* WHO, Bonn.
- WHO (2004) *Meta-analysis of time series studies and panel studies of particulate matter (PM) and ozone (O₃), Eur/04/5042688* WHO Regional Office for Europe, Copenhagen.

8 Climate change, ground level UV and health

(See also section 4.8 of 2001 report)

Graham Bentham, School of Environmental Sciences, University of East Anglia

Summary

- Further epidemiological studies have confirmed the role of UV in skin cancers, and new evidence suggests that UV exposure is a cause of some types of cataract.
- There is growing evidence of important health benefits from sunlight exposure via its effects on vitamin D status.
- The ozone layer is expected to recover by 2050, but climate change may delay that recovery.
- Longer summers and changes in cloud cover may lead to changes in behaviour that are probably more important for personal UV exposure.

Recommendations

- There is a continuing need to maintain health promotion campaigns to limit personal exposure to damaging UV radiation.
- Clearer advice is needed on how the public can maintain adequate vitamin D status in ways that are compatible with the avoidance of risks from exposure to sunlight.

8.1 Introduction

The previous review outlined the ways in which stratospheric ozone depletion was increasing the amounts of biologically active solar ultraviolet radiation reaching the surface. Greenhouse-gas induced climate change may also increase surface level of ultraviolet radiation by mechanisms such as decreases in cloud cover during Spring and Summer. If there were no changes in behaviour increases in UV radiation in the environment would lead to increases in the doses received by the population and any health effects associated with such exposures.

8.2 Health impacts of UV

Apart from an enhanced potential for sunburn, the most important impact would be likely to be an increase in the risk of skin cancers. Another potential effect would be an increase in the incidence of cataracts which are a major cause of blindness. There are also concerns about the ability of UVB radiation to cause suppression of the immune response which could adversely affect the course of some infectious diseases in humans and reduce the effectiveness of vaccinations. On the other hand the evidence was noted that enhanced exposure to ultraviolet radiation might have some health benefits. The most obvious of these is that exposure to sunlight is important for the synthesis of vitamin D which is suspected to have a role in a number of chronic diseases including poor bone health, hypertension, certain auto-immune diseases and some cancers. In the cases of skin cancer and cataracts the evidence was considered strong enough to provide quantitative estimates of the potential health effects of the anticipated increases in surface level of UV radiation.

New studies continue to confirm the crucial role of exposure to sunlight in the aetiology of skin cancers (UNEP, 2003). New animal models have been developed for cutaneous melanoma and for basal cell carcinoma that point to an important role for UVB and especially to exposure early in life.

At the time of the last review some uncertainties remained about the role of sunlight exposure in the formation of cataract which is the leading cause of blindness in the World. New studies from Australia (Neale *et al*, 2003) and from France (Delcourt *et al*, 2000) and a review of 22 published studies (McCarty and Taylor, 2002) all support an association between sun exposure and cataract with animal models (UNEP, 2003) particularly implicating UVB. Knowledge continues to accumulate on the mechanisms by which exposure to UV radiation causes immune suppression but direct evidence on what the implications are for human health remains elusive (UNEP, 2003).

The previous review noted that in addition to the adverse effects of exposure to sunlight there may also be some health benefits although much of the evidence was weak and controversial. In the intervening years this has been an active area of research and a number of well conducted studies have been published supporting the existence of some beneficial effects. Most of these have concentrated on the effects of vitamin D where, for most people, casual exposure to sunlight is the main source. Holick (2004) provides an extensive review of the growing evidence of the importance of sunlight exposure for vitamin D status and role in bone health, the prevention of some cancers, cardiovascular disease and autoimmune diseases such as multiple sclerosis, Type 1 diabetes and rheumatoid arthritis.

8.3 *Changing exposures due to changes in behaviour, climate and ozone layer*

As well as affecting surface levels of ultraviolet radiation by changes in cloud cover it was noted that climate change might have important effects on patterns of behaviour that in turn could have consequences for health. The health effects of solar UV radiation are related to the doses received by the population and these depend on both the UV flux in the environment and patterns of behaviour which bring about exposure. Changes in climate such as increases in sunshine, reductions in precipitation and higher temperatures would all be likely to favour patterns of behaviour involving more outdoor activity, lighter clothing and greater exposure to the sun. At the moment summers in Britain are often too dull and cold to encourage an outdoor lifestyle but a move to sunnier and warmer conditions could have a substantial effect on the UV doses received by the population and the associated health effects. Unfortunately, very little empirical evidence was available on the possible links between climate, outdoor activity and UV exposure in the UK which would allow a quantification of the risks that might be involved.

Since the last review there has been a large volume of published work on trends in stratospheric ozone and associated changes in surface levels of UV radiation. An active research effort has also yielded new information on the potential health impacts of enhanced exposure to UVB. A comprehensive review of this large body of new evidence lies beyond the scope of this report but fortunately major reviews have been published both of ozone depletion and its causes (World Meteorological Organization, 2003) and of its potential impacts on human health (United Nations Environment Programme, 2003).

The WMO review of ozone depletion confirms that a slow recovery of stratospheric ozone is likely as the atmospheric burden of halogens decreases with values reaching pre-Antarctic-ozone-hole value by about mid century. This means that enhanced levels of surface UVB radiation as a result of ozone depletion are likely to persist until approximately 2050. Up to then there will be a period when any effects on human health due to ozone depletion will overlap with those that may be caused by climate change. An important issue on which there is some new data is whether stratospheric cooling related to global warming could delay recovery (McKenzie *et al*, 2003). It has been calculated that a delay of 15-20 years would increase the peak excess incidence of skin cancer in North-West Europe attributable to ozone depletion from 9% above baseline to 15% with the peak year shifting from 2055 to 2065 (UNEP, 2003).

8.4 *Future impacts on health*

Sunburn is perhaps the most obvious health consequence of increased exposure to UV radiation and the previous review noted that particular risks might result from Springtime episodes of ozone depletion acting on a population with largely unacclimatised skin. An interesting new study from an area of Southern Chile periodically affected by the Antarctic ozone hole confirms a large increase in the incidence of sunburn during an episode of ozone depletion that coincided with sunny weather at a weekend (Abarca *et al*, 2002).

The carcinogenic potential of exposure to sunlight may be enhanced by increases in temperature (van der Luen and de Gruijl, 2003) although the evidence is speculative. This study estimates that a 2°C increase in ambient temperature might result in 21% increase in the incidence of skin cancer which is substantially greater than any anticipated effects of ozone depletion alone. However, this is based on extrapolation from the results of experiments on mice and there is, as yet, no direct evidence for humans.

The previous review emphasised that any health effects will be related to the doses of ultraviolet radiation received by the population which will depend on the combination of levels in the environment and patterns of behaviour that bring about exposure. Climate change can be expected to have an effect on both of these. The UKCIP02 climate change scenarios point to larger changes in summer cloud cover than the UKCIP98 scenarios that were the basis of the previous review. The UKCIP02 scenarios for 2080 point to decreases in summer cloud cover for much of England of 9-12% for Medium-Low emissions and 12-15% for Medium-High which, other things being equal, would be associated with increases in UVB radiation of ~5% and ~7% respectively. Previously it was argued that changes in climate such as increases in sunshine, reductions in precipitation and higher temperatures would be likely to favour patterns of behaviour involving more outdoor activity, lighter clothing and greater exposure to the sun.

The likelihood of such changes is supported by evidence from a recently completed study of factors influencing sun exposure of beach users in East Anglia (Horton, 2004). This showed that the length of time spent on the beach and the estimated UVB doses received were strongly positively associated with ambient temperatures whereas the proportion of the body covered by clothing showed a significant negative association. The odds ratio per 1 °C for receipt of a UV dose of >1 minimal erythema dose (MED) was 1.34 (95% C.I. 1.21 to 1.48) which underlines the strong influence of ambient temperature on patterns of behaviour involving potentially risky UV exposure. These findings support the view that that over this century behaviour associated with climate change, rather than ozone depletion, may be the largest determinant of sun exposure and consequent impact on skin cancer (and other health effects) in the UK population (Diffey, 2004).

References

- Abarca, J.F., Casiccia, C.C. and Zamorano, F.D. (2002) Increase in sunburns and photosensitivity disorders at the edge of the Antarctic ozone hole, Southern Chile, 1986-2000. *J.Am.Acad.Dematol.* **46**;2:193-9.
- Delcourt, C., Carriere, I., Ponton-Sanchez, A., Lacroux, A., Covacho, M.J. and Papoz, L. (2000) Light exposure and the risk of cortical, nuclear and sub-capsular cataracts. The Pathologies Oculaires Liees a l'Age (POLA) study. *Archives of Ophthalmology* **118**;3:385-92.
- Diffey, B. (2004) Climate change, ozone depletion and the impact on ultraviolet exposure of human skin. *Physics in Medicine and Biology* **49**;1:R1-R11.
- Holick, M.F. (2004) Sunlight and vitamin D for bone health and prevention of autoimmune diseases, cancers, and cardiovascular disease. *Am.J.Clin.Nutr.* **80**(suppl):1678S-88S.
- Horton B. (2004) An interdisciplinary model of risk from ultraviolet radiation exposure. PhD Thesis. University of East Anglia. Norwich, 2004.
- McCarty, C.A. and Taylor, H.R. (2002) A review of the epidemiological evidence linking ultraviolet radiation and cataracts. *Developments in Ophthalmology* **35**:21-31.
- McKenzie, R.L., Bjorn, L.O., Bais, A. and Ilyasad, M. (2003) Changes in biologically active ultraviolet radiation reaching the Earth's surface. *Photochem.Photobiol.Sci.* **2**;1:5-15.
- Neale, R.E., Purdie, J.L., Hirst, L.W. and Green, A.C. (2003) Sun exposure as a risk factor for nuclear cataract. *Epidemiology* **14**;6:707-12.
- United Nations Environment Programme. Environmental Effects of Ozone Depletion and its Interactions with Climate Change: 2002 Assessment. Nairobi, 2003.
- Van der Leun, J.C. and de Gruijl, F.R. (2002) Climate change and skin cancer. *Photochem.Photobiol.* **1**:324-6.
- World Meteorological Organization. Scientific Assessment of Ozone Depletion: 2002, Global Ozone Research and Monitoring Project—Report No. 47, Geneva, 2003.